



Pulsed Electric Field Technology for Innovative Food Processing

Haruka Tanaka*

Department of Food Technology and Nutritional Science, Institute of Food Innovation, Osaka, Japan

DESCRIPTION

Pulsed Electric Field (PEF) technology is an emerging food-processing method that uses short-duration electrical pulses to modify biological materials and control microorganisms. It is particularly relevant to liquid and semi-liquid foods because electrical energy can be delivered rapidly through a continuous treatment chamber. PEF has attracted scientific interest because it can support microbial inactivation, improve extraction efficiency and modify plant tissues while reducing dependence on prolonged thermal exposure.

The central mechanism of PEF is electroporation. When a sufficiently strong electric field is applied to biological cells, changes occur in the structure of cellular membranes. Temporary pores may form when treatment intensity is moderate, whereas stronger exposure can cause irreversible membrane disruption. The extent of electroporation depends on electric-field strength, pulse duration, number of pulses, temperature, conductivity and the characteristics of the cells being treated.

It has been investigated extensively for fruit and vegetable juices. Microbial inactivation can be achieved when appropriate electrical conditions are selected, potentially allowing processors to reduce the severity of thermal treatment. This can be useful for products in which color, aroma and heat-sensitive nutrients are commercially important. The actual effectiveness, however, depends on the food matrix and target microorganisms, so process validation remains essential.

Another important application is extraction enhancement. Plant cells contain valuable compounds within structures that can restrict their release during conventional processing. PEF-induced electroporation can increase cell permeability and facilitate the movement of intracellular components. This principle has been explored for improving juice yield, recovering pigments and extracting phenolic compounds and other functional constituents.

It can also act as a pretreatment before drying or pressing. Modification of plant tissue may improve mass transfer and

reduce the resistance encountered during subsequent processing. In some applications, this could shorten drying times or increase extraction yield. Such process intensification is particularly interesting for food manufacturers attempting to reduce energy use and improve raw-material utilization.

The electrical conductivity of a product is an important processing parameter. Foods with greater conductivity allow electrical current to pass more readily, whereas low-conductivity products may require different operating conditions. Conductivity can change during processing as temperature increases or as components are released from food tissues. Consequently, PEF systems must be designed to accommodate changes in electrical behavior.

Equipment design is another important consideration. Industrial PEF systems commonly operate in continuous flow, allowing products to pass between electrodes within a controlled treatment chamber. Electrode geometry, insulation, flow rate, electric-field distribution and treatment temperature influence performance. The design must also minimize unwanted electrical phenomena and maintain hygienic processing conditions.

PEF has several potential advantages over conventional processing. Very short treatment periods can reduce thermal exposure, while continuous operation may support high throughput. The technology can also be combined with other preservation methods. For example, it may be used as a pretreatment followed by mild heating or refrigeration. Such hurdle approaches can improve microbial control without relying excessively on one processing method.

Despite its promise, PEF has limitations. Highly viscous foods, products with large particles and materials with heterogeneous electrical properties can be difficult to process uniformly. Scale-up from laboratory equipment to industrial systems can also introduce challenges related to fluid dynamics and electric-field distribution. Equipment cost and energy management must be considered when assessing commercial feasibility. Therefore, further research is needed to optimize processing parameters, improve equipment efficiency, and expand PEF.

Correspondence to: Haruka Tanaka, Department of Food Technology and Nutritional Science, Institute of Food Innovation, Osaka, Japan. Email: haruka.tanaka@612.com

Received: 21-Nov-25, Manuscript No. JFPT-25-32875; **Editor assigned:** 24-Nov-2025, Pre QC No. JFPT-25-32875 (PQ); **Reviewed:** 08-Dec-2025, QC No. JFPT-25-32875; **Revised:** 15-Dec-2025, Manuscript No. JFPT-25-32875 (R); **Published:** 22-Dec-2025, DOI: 10.35248/2157-7110.25.16.1189

Citation: Tanaka H (2025). Pulsed Electric Field Technology for Innovative Food Processing. J Food Process Technol.16:1189.

Copyright: © 2025 Tanaka H. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

Food safety remains the primary requirement. Treatment conditions must demonstrate adequate reduction of relevant microorganisms under realistic commercial conditions. Microbial resistance can vary with physiological state, growth environment, acidity, water activity and food composition. Therefore, laboratory measurements should be supported by product-specific validation.

Future research may improve through better treatment chambers, real-time monitoring, predictive modeling and integration with other processing technologies. This may

ultimately contribute not only to preservation but also to extraction, drying and process intensification.

In conclusion, pulsed electric field technology therefore represents a multifunctional approach to modern food processing. Its ability to interact directly with biological structures provides opportunities for safer processing, improved extraction and reduced processing intensity. Continued technological development will be important for transforming promising laboratory applications into reliable industrial operations.