



Mechanisms of Genomic Instability in Human Cancer

Adrian Keller*

Department of Cancer Genomics, Institute of Molecular Oncology, Ludwig Maximilian University of Munich, Munich, Germany

DESCRIPTION

Genomic instability refers to an increased tendency of the genome to acquire genetic alterations during cell division. It is considered one of the fundamental characteristics of cancer and plays a central role in tumor initiation, progression, and therapeutic resistance. Under normal physiological conditions, cells possess highly efficient mechanisms that maintain genomic integrity by accurately replicating Deoxyribonucleic Acid (DNA), genetic material, and ensuring the faithful segregation of chromosomes during mitosis. When these protective mechanisms become impaired, mutations accumulate throughout the genome, leading to chromosomal abnormalities, gene amplifications, deletions, translocations, and other structural alterations that promote malignant transformation. The continuous accumulation of genetic changes provides cancer cells with evolutionary advantages that facilitate uncontrolled growth, invasion, and adaptation to changing environmental conditions.

Genomic stability is preserved through a complex network of DNA repair pathways, cell cycle checkpoints, and chromosome maintenance systems. DNA is continuously exposed to endogenous and exogenous sources of damage, including reactive oxygen species generated during cellular metabolism, ultraviolet radiation, ionizing radiation, environmental chemicals, and replication errors. To counteract these insults, cells utilize multiple DNA repair mechanisms such as base excision repair, nucleotide excision repair, mismatch repair, homologous recombination, and non-homologous end joining. These pathways detect and repair damaged DNA before mutations become permanently incorporated into the genome. Failure of one or more repair systems significantly increases mutation frequency and contributes to genomic instability.

Several forms of genomic instability have been identified in human cancers. Chromosomal instability involves gains, losses, or rearrangements of whole chromosomes or chromosomal segments, resulting in aneuploidy and structural abnormalities.

Microsatellite instability arises from defects in the DNA mismatch repair system, causing alterations in short repetitive DNA sequences distributed throughout the genome. Elevated mutation rates resulting from defective DNA repair represent another form of genomic instability and contribute to the accumulation of numerous point mutations across cancer genomes. These different types of instability may coexist within the same tumor, generating substantial genetic diversity among cancer cells and influencing disease progression and therapeutic response.

Numerous molecular abnormalities contribute to the development of genomic instability. Mutations affecting tumor suppressor genes involved in DNA repair and cell cycle regulation impair the ability of cells to recognize and repair damaged DNA. Defective chromosome segregation during mitosis may result in abnormal chromosome numbers and structural rearrangements. Telomere shortening and dysfunction also contribute to genomic instability by promoting chromosome fusion and breakage during successive cell divisions. Oxidative stress, chronic inflammation, and persistent exposure to environmental mutagens further increase DNA damage and overwhelm cellular repair mechanisms. Together, these alterations create an environment in which genetic abnormalities accumulate rapidly, accelerating the evolution of malignant cells.

In conclusion, genomic instability is a defining feature of cancer that drives the accumulation of genetic alterations responsible for tumor initiation, progression, metastasis, and therapeutic resistance. Defects in DNA repair, chromosome maintenance, and cell cycle regulation collectively contribute to this process and create opportunities for malignant evolution. Continued advances in molecular diagnostics and precision medicine are improving the ability to detect genomic instability, identify high-risk individuals, and develop targeted therapeutic strategies. A deeper understanding of the mechanisms underlying genomic instability will continue to enhance cancer prevention, early diagnosis, and individualized treatment, ultimately improving patient outcomes and advancing the field of oncology.

Correspondence to: Adrian Keller, Department of Cancer Genomics, Institute of Molecular Oncology, Ludwig Maximilian University of Munich, Munich, Germany, E-mail: adrian.keller@lmu.de

Received: 22-Apr-2026, Manuscript No. JCM-26-32081; **Editor Assigned:** 24-Apr-2026, PreQC No. JCM-26-32081 (PQ); **Reviewed:** 08-May-2026, QC No. JCM-26-32081; **Revised:** 15-May-2026, Manuscript No. JCM-26-32081(R); **Published:** 22-May-2026, DOI: 10.35248/2157-2518.26. S59.003

Citation: Keller A (2026). Mechanisms of Genomic Instability in Human Cancer. S59:003.

Copyright: ©2026 Keller A. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.