



## Applications of Biotechnology in Food Sciences

Franco Ferrari\*

Department of Food Technology, Illinois University, Chicago, USA

### ABOUT THE STUDY

Food technology is a branch of food science that deals with the production, preservation, quality control and research and development of the food products. Early scientific research into food technology concentrated on food preservation. Food biotechnology is an term covering a vast variety of processes for using living organisms similar as shops, creatures, microbes, or any part of these organisms to develop new or advanced food products. It includes the newer forms of food biotechnology that offer a briskly and more precise means to develop food products. Food biotechnology is the operation of ultramodern biotechnological ways to the manufacture and processing of food products as well as food constituents and food complements. Food biotechnology, including inheritable revision of foods and micro-organisms, nutritive genomics and the development of functional foods. Turmoil improves the delectability and adequacy of raw accoutrements by producing flavour and aroma factors and modifying food texture [1].

Old Aged food biotechnology ways include conventional crossbreeding, which refers to the arbitrary recombination of genes through sexual reduplication leading to a new organism with bettered traits. Crossbred shops, for case, may bear several generations to achieve a particular particularity due to the randomness of gene transfer. Exemplifications of similar traits are bettered crop yield, aesthetic rates, increased forbearance to physical stress similar as cold temperatures, and increased resistance to complaint and insects.

New Modern food biotechnology ways include the joining of two pieces of DNA from different organisms leading to a single piece of DNA. Individual "specific" genes are transferred from one organism to another in order to ameliorate the nutrient situations of a food, for illustration, similar as fortifying a fruit or vegetable [2]. Ultramodern ways are important briskly and more precise. It's possible to snappily transfer a specific gene of interest rather than staying on the arbitrary shuffling of genes over several generations.

Exemplifications of foods developed through biotechnology to increase the situations of nutrients or to address a health concern include canvases, similar as canola, in which the situations of nutritionally essential adipose acids are increased, kinds of wheat that don't contain gluten, and potatoes (protein), kiwi (resveratrol), and lettuce (iron) [3].

Incentive strains used for wine conflation are able of malolactic turmoil. Wine conflation consists of two way, Primary turmoil results in conversion of glucose into alcohol using incentive, Secondary turmoil uses bacteria and its product is lactic acid and this causes the rise in position of acidity. To overcome this problem different strategies are used which are expensive. This problem was answered through insertion of malolactic gene (*Lactobacillus delbrueckii*) in artificial incentive strain. This gene lowers the malate conversion hence lowering acidity position of wine [4].

Using biotechnology in the growth and product of fruits and vegetables has enabled scientists to change the way they grow. Typically fruits and vegetables continue to grow after harvesting; they must be rushed to request and vended snappily while they're fresh. Genetically modified yield can be gathered when ripe, and the growing process stops, giving them a longer shelf life. These inheritable variations also increase a factory's resistance to complaint, pests, germicides, dressings and indeed extreme rainfall conditions. Inheritable engineering has also altered a factory's nutritive makeup, making it richer in certain vitamins or minerals [5,6].

Disadvantages of harvesting thousands of genetically modified shops, scientists still don't have a clear understanding of how these inheritable changes affect the mortal body long term, primarily because these food shops have only been available since the early 1990s. Monsanto, the largest GM seed inventor in the world and the association that provides a maturity of the exploration results to the FDA, says GM seed is inoffensive to humans. Still, some scientists have plant that GM shops have indeed altered the life span, complaint process and cognitive capacities of insects that feed on these shops.

**Correspondence to:** Franco Ferrari, Department of Food Technology, Illinois University, Chicago, USA, E-mail: fferrari@iit.edu

**Received:** 01-Apr-2022, Manuscript No. JFPT-22-16646; **Editor assigned:** 04-Apr-2022, PreQC No. JFPT-22-16646 (PQ); **Reviewed:** 18-Apr-2022, QC No JFPT-22-16646; **Revised:** 25-Apr-2022, Manuscript No. JFPT-22-16646 (R); **Published:** 02-May-2022. DOI: 10.35248/2157-7110.22.13.925

**Citation:** Ferrari F (2022) Applications of Biotechnology in Food Sciences. J Food Process Technol. 13: 925.

**Copyright:** © 2022 Ferrari F. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

## References

1. Williams JB, Napoli JL. Metabolism of retinoic acid and retinol during differentiation of F9 embryonal carcinoma cells. *Proc Natl Acad Sci.* 1985;82(14):4658-4662.
2. Lee TF, Mak KM, Rackovsky O, Lin YL, Kwong AJ, Loke JC, et al. Downregulation of hepatic stellate cell activation by retinol and palmitate mediated by adipose differentiation-related protein (ADRP). *J Cell Physiol.* 2010;223(3):648-657.
3. Cammas L, Romand R, Fraulob V, Mura C, Dolle P. Expression of the murine retinol dehydrogenase 10 (Rdh10) gene correlates with many sites of retinoid signalling during embryogenesis and organ differentiation. *Dev Dyn.* 2007;236(10):2899-2908.
4. Janick-Buckner D, Barua AB, Olson JA. Induction of HL-60 cell differentiation by water-soluble and nitrogen-containing conjugates of retinoic acid and retinol. *FASEB J.* 1991;5(3):320-325.
5. Marx J, Naudé H, Pretorius E. The effects of hypo-and hypervitaminosis a and Its involvement in foetal nervous system development and post-natal sensorimotor functioning–A Review. *Br J Dev Disabil.* 2006;52(102):47-64.
6. Zile MH. Vitamin A and embryonic development: an overview. *J Nutr.* 1998;128(2):455-458.