

Assessment of Major and Trace Elements in Aquatic Macrophytes, Soils and Bottom Sediments Collected Along Different Water Objects in the Black Sea Coastal Zone by Using Neutron Activation Analysis

Nekhoroshkov PS^{1*}, Kravtsova AV², Kamnev AN³, Duliu O⁴ Bunkova OM³, Frontasyeva MV¹ and Yermakov IP²

¹Frank Laboratory of Neutron Physics, Joint Institute for Nuclear Research, Russia

²Faculty of Biology, Lomonosov Moscow State University, Moscow, Russia

³Faculty of Soil Science, Lomonosov Moscow State University, Moscow, Russia

⁴Department of the Structure of Matter, Faculty of Physics, University of Bucharest, Romania

Abstract

The levels and compartmentalization of Na, Mg, Al, Cl, K, Ca, Sc, Ti, V, Cr, Mn, Fe, Co, Ni, Zn, As, Se, Br, Rb, Sr, Mo, Sb, I, Cs, Ba, La, Ce, Sm, Eu, Tb, Hf, Ta, Au, Th, and U in *Phragmites australis* *Carex conescens* L and *Cladophora sericea* from the Caucasian coast of the Black Sea Anapa recreational region was investigated by Neutron Activation Analysis. The study touches upon subject of the sediment-to-plant and root-to-leaf elemental transfer as well as of the influence of anthropogenic pollution on wetland ecosystems in zone of resort. The content of the majority of considered elements was found higher in the belowground organs of *P. australis* than in the aboveground tissues while a reverse regularity was evidenced for *C. conescens*. The levels of elements decrease from bottom sediments to aquatic plants with the notable exception of the halogens Cl, Br and I that presented 5 to 100 fold higher content in plants than in sediments. The increased levels of As, Mo, and Sb in some soil and sediment samples most probably indicate the anthropogenic pollution. It recommends them for a continuous monitoring of the same area.

Keywords: Major and trace elements; Neutron activation analysis; Black Sea; *Phragmites australis*; *Carex conescens*; *Cladophora sericea*

Introduction

The aquatic macrophytes are widely used for assessing the environmental situation in fresh as well as seawater [1-10]. As the accumulation of trace metals in organisms depends on the concentration of pollutants in water and sediments as well as on exposure time, a tissue analysis of aquatic macrophytes may provide cumulative evaluation of exposure [11-13]. The concentrations of chemical elements in aquatic plants can be more than 100,000 times higher than in the associated water [14]. This accumulation ability of certain macrophytes is used for monitoring purposes in relatively clean and recreation zones, where low level of contamination might be difficult to detect [15]. Our previous investigations in the Black Sea [16-18] evidenced for the increased elemental concentrations in marine algae in more polluted waters reflecting their great potential for biomonitoring of water quality. They proved not only the existence of a certain degree of anthropogenic contamination but also the suitability of aquatic plants for biomonitoring of trace elements. To extend our studies regarding the elemental content of more than 35 elements for the territory of an important but poor investigated recreation zone of the Caucasian coast of the Black Sea, the aquatic macrophytes *Phragmites australis*, *Carex conescens* L as well as the green algae *Cladophora sericea* (Hudson) Kutzing were used. *P. australis* is one of the most distributed macrophytes in aquatic ecosystems, and numerous studies showed its capacity of trace element bioaccumulation [19-23]. Thus Duman reported that the roots of *Phragmites australis* from fresh water Lake Sapanca in Turkey were found to be good accumulators of Cu, Mn, Ni, Zn. The studies of in the estuaries of Italian rivers affected by municipal wastewaters and agricultural activities showed a good correlation of Al, As, Cr, Cu, Mn, Ni and Zn in *P. australis* with the elemental content in corresponding sediments and water. Also a strong positive correlation between the concentrations of Al, As, Co, Cr, Cu, Fe, Mn, Ni, Se, Sr and Zn in the sediments and all organs (rhizome, stem and leaf) of *P. australis* sampled from the Tisza River in Serbia was found by [23]. The investigations in the constructed wetland in North Italy [24] and in the Hokersar wetland, Ramsar site of Kashmir Himalaya, India [25] showed that *P. australis* is appropriate species for phytoextraction

and phytoremediation of the environment. Analysis of the elemental composition of *P. australis*, collected in the Anapa region in 2013-2014, showed that the concentration ratios with the absolute value that is greater than 1 (pointing to the pollution of the area) are determined only for As. Maximal values of biological absorption coefficients were found for the As, Fe, K, Mn, Zn in roots. The data of using the species of *Carex* (sedges) in biomonitoring purposes are scarce in comparison with *Phragmites*. Horovitz [26] reported the content of Ag, Co, Cr, Cs, Fe, Rb, Sc, Th, Zn and in *Carex pendula* sampled in botanical garden in Germany. Pederson and Harper [27] studied the chemical composition of some major forage plants of mountain summer ranges of southeastern Utah, USA, reported the content of K, Ca and Mg in *Carex geyeri*. Ohlson [28] studied the content of Al, Ca, Cu, Fe, K, Na, Mg, Mn, Mo, Zn in eleven plants from the mires of central and north Sweden found that the largest variation in elemental concentration of roots and leaves was observed in *Carex rostra*. He also reported that the concentration of K in tissues of *Carex* species was highly correlated with its concentration in the substrate. The species of green algae of genus *Cladophora* has frequently been suggested as a suitable organism to monitor water contamination and its practical use in monitoring river, lake and sea pollution has been reported from a range of countries [29-34]. Thus Whitton reported that there were highly significant correlation between Cu, Fe, Zn content in *Cladophora glomerata* from rivers and streams in Northern England and water. The similar results

*Corresponding author: Nekhoroshkov PS, Frank Laboratory of Neutron Physics, Joint Institute for Nuclear Research, Russia, Tel: +74962163834; E-mail: p.nekhoroshkov@gmail.com

Received November 30, 2016; Accepted December 30, 2016; Published January 03, 2017

Citation: Nekhoroshkov PS, Kravtsova AV, Kamnev AN, Bunkova O (2017) Assessment of Major and Trace Elements in Aquatic Macrophytes, Soils and Bottom Sediments Collected Along Different Water Objects in the Black Sea Coastal Zone by Using Neutron Activation Analysis. Mod Chem Appl 5: 208. doi: 10.4172/2329-6798.1000208

Copyright: © 2017 Nekhoroshkov PS, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

were reported by for Cr, Ni and V determined in *Cladophora glomerata* from refinery sewage lagoon (Bratislava). Levkov and Krstic found that the levels of Co, Cu, Fe, Mn and Zn in *Cladophora glomerata* reflected their load in the River Vardar, Macedonia, and recommended it as a precise biomonitoring tool for determination and quantification of heavy metal pollution in this river. In the distribution patterns of Ca, Cu, K, Mg, Mn, Na, Ni, Zn and in the green algae *Cladophora* sp. from the Southern Baltic is assessed. The study concluded that *Cladophora* sp. can be used the most successfully as biomonitor of Cu and Zn content in the Baltic Sea because of its ability to accumulate metal contaminants from seawater, tolerance to metals, simple morphology and adequate tissue for analysis. The preliminary study of elemental composition of *Cladophora sericea*, collected in the Anapa region in 2013, showed that the plant to soil ratios greater than one and pointing towards a possible contamination process were detected only for As, and Sr. For our study, we have chosen three types of phototrophic macrophytes as ones of the most convenient organisms-biomonitor. Moreover, they occur in different ecological conditions and are the first ones that take the fall of the coastal pollution runoff accordingly, we have investigated the hydrophyte filamentous marine green alga *Cladophora sericea* (Hudson) Kutzing, helophyte *Phragmites australis* (Cav) Trin. Ex Steud as well as the hygrophyte *Carex conescens* *Cladophora sericea* lives in shallow sandy areas of the Black Sea; absorb major and trace elements by all surface of their body, do not have root system, *Phragmites australis* lives along the coastal zones of rivers and seas. The well-developed root system makes more than 80% of the total biomass. Plants absorb major and trace elements from soil, sediment and water by additional roots. *Carex conescens* grows on the banks of the rivers. Unlike *Phragmites australis* is plant has a small root system so it absorbs major and trace elements only from the soil. The main goals of the study consist of: quantifying the content of a wide range of major as well as trace elements in *Phragmites australis*, *Carex conescens*, *Cladophora sericea* and corresponding soil and bottom sediments samples assessing the elemental content in different parts of plants (leaves and roots) quantifying the element mobility from sediment to organs, as well as within the plant, providing new data on the geochemistry of sediments and soil from the Anapa region quantifying the level of the anthropogenic pollution of the study area. The results of this project will be further presented and discussed.

Materials and Methods

Study area and sampling

A resort city of Anapa (Krasnodar region) located on the Caucasian

coast of the Black Sea is characterized by humid subtropical climate and long sandy beach. The Anapka river crossing the territory of the town connects Anapa reed beds with the Black Sea [35]. The investigated area (Figure 1; Table 1) includes the municipal waste dump at the Krasnyi hutur and some reservoirs, i.e., a lake, a river and reed beds at the foot of the hill and below the dump situated on the highland. These water bodies form an indivisible watershed of the river Anapka which estuary occupies the main city beach within the city recreation zone. The solid waste city dump of Anapa is located near the Krasnyi hutur, 4.65 km from the Black Sea. During 10 years, the total area of the dump increased from 9 to 26 hectares in 2013. There is a lake located in the distance of 0.62 mile downhill from the dump at village Krasnyi hutur 7). The next sampling point is Anapa reed beds this marshland is situated at the hollow, where the Kotloma and Kumatyr Rivers get its confluence, not far from the Anapa station. The length of the Anapka reed beds is 1 mile long. The station 2b is situated in old bed of the Anapka river. The mouth of the Anapka river is located at the main city beach. The samples of vegetation (live and dead leaves and roots of *Phragmites* and *Carex*, algae *Cladophora*) (n=35) and the corresponding soil (n=40) and bottom sediment (BS) (n=15) were collected at 7 sites




Figure 1: Sampling sites in the studied area near Anapa city (Anapa transect or cross-section).

Sampling point	Latitude (N)	Longitude (E)	Type	Summary description
City dump	44°57'31.76"	37°21'51.01"	Soil	Dump without vegetation
Lake near Krasnyi Hutur (village)	44°56'51.52"	37°20'41.70"	Sediments Plants	Waste liquid disposal, polluted runoff
Anapa reed beds	44°55'35.97"	37°19'47.57"	Soil Sediments Plants	Traffic, Gas station
Anapka river	44°54'35.84"	37°19'45.33"	Soil Sediments Plants	Traffic
Old bed of Anapka river	44°54'10.27"	37°19'10.18"	Soil Sediments Plants	Beach, objects of recreation
Mouth of Anapka river	44°54'21.59"	37°19'06.89"	Soil Sediments Plants	Beach
Anapa Bay	44°54'11.74"	37°19'06.87"	Sediments Plants	Beach, marine traffic

Table 1: The location of sampling points as well as the type of collected material.

along the transect located near Anapa city in summer of 2013 and 2014. The sampling sites are shown in Figure 1 while Table 2 presents more details regarding the sampling points location as well as a summary description of each category of samples. The samples of soil and bottom sediments were collected according to GOST (state standard) [36,37]. The sampling process ensures compliance with the two requirements




Figure 2: The diagram of the elemental content in soils and sediments normalized on UCC for 7 and 2c stations.

the amount of the sampled material to be enough for the analysis, and in the case of soils or sediments, their texture should be an average one over the studied media. Both soil and bottom sediments samples were air dry in a warm and ventilated room until constant weight, then milled and sieved through 0.04 mm mesh. Special attention was given to the preparation of the plant material samples, which was dried at 105°C until constant weight, ground in an agate mortar. There were no further chemical preparations of the samples, which excluded the errors due to the reagents.

Neutron activation analysis

Elemental analysis of the samples was carried out by INAA at the reactor IBR-2 of the Frank Laboratory of Neutron Physics (FLNP) of the Joint Institute for Nuclear Research (JINR), Dubna, Russia. The analytical procedures and the basic characteristics of the employed experimental facility are described in detail elsewhere [38]. The samples of about 0.3 g were packed in polyethylene bags for short-term irradiation and in aluminum cups for long-term irradiation. To determine the short-lived isotopes of Mg, Al, Cl, Ca, Ti, V, Mn and I the samples were irradiated

El.	SRM 1632c		SRM 433		SRM 667	
	Certified	Determined	Certified	Determined	Certified	Determined
Na	299 ± 5	300 ± 8	13500 ± 4050	13150 ± 160	-	-
Mg	384 ± 32	362 ± 15	11500 ± 230	11430 ± 120	-	-
Al	9150 ± 137	9350 ± 187	78200 ± 782	77980 ± 890	-	-
Cl	1139 ± 41	1120 ± 36	-	-	-	-
K	1100 ± 33	1100 ± 201	16600 ± 2224	16300 ± 250	-	-
Ca	1450 ± 290	1430 ± 130	-	-	-	-
Sc	2.9 ± 0.03	2.91 ± 0.07	14.6 ± 4.38	15.1 ± 0.15	13.7 ± 0.7	12.3 ± 0.24
Ti	517 ± 32	511 ± 21	-	-	-	-
V	23.7 ± 0.52	25.3 ± 0.78	160 ± 2.08	152 ± 11	-	-
Cr	8.24	13.7	136 ± 1	136 ± 4	178 ± 16	172 ± 8.5
Mn	13 ± 0.52	13.2 ± 0.46	316 ± 3.16	313 ± 5	920 ± 40	924 ± 18
Fe	7350 ± 110	7350 ± 250	40800 ± 408	40805 ± 1673	44800 ± 986	39926 ± 1200
Co	3.48 ± 0.2	3.91 ± 0.24	39.4 ± 0.4	39.4 ± 2.9	23 ± 1.3	19 ± 0.2
Ni	9.32 ± 0.51	10.5 ± 3.2	39.4 ± 0.39	39.4 ± 0.14	128 ± 8.96	23 ± 1
Zn	12.1 ± 1.29	11.2 ± 1.8	101 ± 1	101 ± 3	175 ± 13	148 ± 3
As	6.18 ± 0.27	6.25 ± 0.35	18.9 ± 0.2	18.9 ± 0.4	17.1 ± 5.13	17.5 ± 4
Se	1.33 ± 0.07	1.33 ± 0.09	0.78 ± 0.03	0.72 ± 0.3	1.59 ± 0.08	1.49 ± 0.1
Br	18.7 ± 0.39	17.9 ± 0.5	67 ± 7.97	70 ± 5	99.7 ± 2.5	99.7 ± 2.7
Rb	7.5 ± 0.3	7.5 ± 1.3	99.9 ± 8.49	102 ± 14	-	-
Sr	63.8 ± 1.4	63.4 ± 5.3	302 ± 3	302 ± 20	224.5 ± 67.3	200 ± 10
Mo	0.8 ± 0.24	0.79 ± 0.28	-	-	-	-
Sb	0.46 ± 0.03	0.46 ± 0.04	1.96 ± 0.04	1.96 ± 0.06	0.96 ± 0.05	0.74 ± 0.04
I*	-	-	-	-	-	-
Cs	0.59 ± 0.01	0.59 ± 0.02	6.4 ± 0.26	6.2 ± 0.06	7.8 ± 0.7	6.7 ± 0.08
Ba	41 ± 2	41 ± 3	268 ± 19	268 ± 12	-	-
La	-	-	33.7 ± 1.61	31 ± 5	27.8 ± 1	27.8 ± 1.1
Ce	11.9 ± 0.2	17.5 ± 3.6	64.5 ± 19.4	73.9 ± 1.5	56.7 ± 2	57 ± 3
Sm	1.08 ± 0.03	1.08 ± 0.04	-	-	4.66 ± 0.2	4.25 ± 0.5
Eu	0.12 ± 0.003	0.32 ± 0.1	1.18 ± 0.35	2.42 ± 0.04	1 ± 0.01	1.0 ± 0.1
Tb	-	-	0.696 ± 0.2	0.7 ± 0.03	0.68 ± 0.02	0.60 ± 0.02
Hf	0.59 ± 0.01	0.59 ± 0.05	3.66 ± 1.1	4.41 ± 0.08	-	-
Ta	-	-	1.03 ± 0.31	1.00 ± 0.02	0.88 ± 0.02	0.88 ± 0.02
Au	-	-	-	-	0.017 ± 0.005	0.017 ± 0.005
Th	1.4 ± 0.03	1.4 ± 0.04	9.8 ± 0.3	9.8 ± 0.3	10 ± 0.5	9.14 ± 0.09
U	0.51 ± 0.01	0.51 ± 0.02	2.45 ± 0.2	2.23 ± 0.2	2.26 ± 0.15	2.29 ± 0.3

Table 2: The NAA data and certified values of reference materials (mean ± error, in µg g⁻¹ dry weight).

(I concentration was determined using SRM 1515 (apple leaves): certified value 0.3±0.09; determined value 0.26±0.12.)

for 3 min in the reactor channel with a neutron flux density of $1.3 \cdot 10^{12}$ n/(cm² s). Gamma spectra of induced activity were measured for 12-15 min after 20 min of decay. The elemental contents of the long-lived isotopes of Na, K, Sc, Cr, Fe, Co, Ni, Zn, As, Se, Br, Rb, Sr, Mo, Sb, Cs, Ba, La, Ce, Sm, Eu, Tb, Hf, Ta, Au, Th, and U were determined using epithermal neutrons in a cadmium-screened irradiation channel with a neutron flux density of $1.6 \cdot 10^{12}$ n/(cm² s). Samples were irradiated for 90 h, repacked and then measured twice after 4–5 d of decay during 30 min and after 20 days of decay during 1.5 h. To process gamma spectra of induced activity and to calculate concentrations of elements in the samples, software developed at FLNP, JINR was used [39]. The uncertainties in the determined concentrations were in the range of 5-15%, and of 30% or more for those elements which concentrations in the samples were at the detection limit. Quality control was provided by using National Institute for Standard and Technology (NIS) reference materials (SRM): NIST 1632c (trace elements in coal), NIST 433 (marine sediment), NIST 667 (estuarine sediment) as well as NIST 1515 (apple leaves) irradiated in the same conditions together with

the samples under investigation. The NAA data and certified values of reference materials are given in concentration were determined using SRM 1515 (apple leaves): certified value 0.3 ± 0.09 ; determined value 0.26 ± 0.12 .

Data analysis

To unify the major and trace composition of each plant we used the Reference Plant (RP) contents [40] as normalizing factors. In this way, it was possible to compare the distribution of the considered elements in all species of plants chosen for the present study. A similar approach we used in the case of soils and sediments by considering the Upper Continental Crust (UCC) [41] as reference average rock. Therefore, all data regarding the elemental composition of the Anapa soils and sediments samples were normalized to the corresponding content of the UCC. The accurate data on concentrations with the wide number of elements for “average sediment” are presented in UCC. The normalized on UCC data of concentrations in soils and sediments were used for comparison the levels of elements between different stations. The levels

Element	Station 7		Station 2c		UCC ¹	SNC ²
	Soils	BS	Soils	BS		
Na	4200 ± 700	4300 ± 700	8000 ± 2000	7400 ± 2200	24259	–
Mg	20000 ± 6300	10200 ± 5600	5300 ± 3000	5000 ± 3400	14957	–
Al	60000 ± 7000	38000 ± 600	23000 ± 2000	20000 ± 400	81505	–
Cl	260 ± 160	800 ± 300	430 ± 150	240 ± 70	370	–
K	16000 ± 3000	11200 ± 600	8300 ± 1400	8900 ± 2600	23244	–
Ca	44000 ± 6300	73000 ± 11000	68200 ± 10400	75300 ± 12100	25658	–
Sc	12.1 ± 3.5	7.8 ± 0.2	1.48 ± 0.30	1.46 ± 0.01	14	–
Ti	3500 ± 400	2400 ± 300	600 ± 180	450 ± 70	3897	5030
V	136 ± 18	88 ± 5	14 ± 6	11.5 ± 1.2	97	126
Cr	89 ± 25	86 ± 38	12 ± 4	9 ± 2	92	109
Mn	704 ± 116	463 ± 50	210 ± 50	180 ± 13	774	930
Fe	33500 ± 8600	34800 ± 8800	4900 ± 600	4800 ± 100	39176	–
Co	17.7 ± 5.1	15.0 ± 0.1	2.0 ± 0.3	1.9 ± 0.3	17.3	–
Ni	58.2 ± 19.2	53.1 ± 7.8	5.3 ± 1.7	5.5 ± 0.4	47	47
Zn	86.3 ± 11.4	112 ± 51	19 ± 10	14 ± 3	67	106
As	14.8 ± 2.5	17.6 ± 5.1	6.7 ± 0.7	6.5 ± 0.4	4.8	–
Se	1.1 ± 0.9	1.6 ± 1.5	0.20 ± 0.15	0.2 ± 0.1	0.09	–
Br	18.6 ± 4.9	17.3 ± 1.1	3.5 ± 0.7	2.7 ± 0.6	1.6	–
Rb	83.3 ± 21.9	53.8 ± 3.3	27.9 ± 4.5	25.9 ± 0.1	84	–
Sr	370 ± 250	438 ± 49	470 ± 170	500 ± 4	320	216
Mo	6.2 ± 5.3	14.2 ± 14.1	1.0 ± 1.0	1.0 ± 0.1	1.1	–
Sb	1.7 ± 0.4	2.4 ± 0.9	0.20 ± 0.03	0.17 ± 0.01	0.4	–
I	12.3 ± 2.7	16.5 ± 3.2	2.1 ± 0.7	4.5 ± 0.9	1.4	–
Cs	5.5 ± 1.7	3.3 ± 0.04	0.40 ± 0.07	0.40 ± 0.01	4.9	–
Ba	530 ± 170	524 ± 84	225 ± 60	200 ± 41	624	720
La	36.5 ± 18.6	66.9 ± 55.3	10.8 ± 8.2	10.8 ± 6.4	31	–
Ce	41.9 ± 24.5	39.6 ± 6.8	11.5 ± 2.7	13.7 ± 0.4	63	–
Sm	5.6 ± 1.8	9.8 ± 7.4	1.2 ± 0.4	1.3 ± 0.3	4.7	–
Eu	1.7 ± 0.6	0.8 ± 0.9	0.3 ± 0.1	0.30 ± 0.01	1	–
Tb	0.6 ± 0.2	0.6 ± 0.04	0.14 ± 0.03	0.140 ± 0.004	0.7	–
Hf	6.1 ± 1.9	4.3 ± 1.1	1.2 ± 0.3	1.1 ± 0.2	5.3	–
Ta	0.6 ± 0.2	0.5 ± 0.1	0.10 ± 0.03	0.09 ± 0.01	0.9	–
Au	0.01 ± 0.001	0.01 ± 0.001	0.002 ± 0.002	0.002 ± 0.001	1.5	–
Th	8.6 ± 2.1	7.6 ± 0.3	1.4 ± 0.3	1.27 ± 0.02	10.5	–
U	4.3 ± 6.5	2.2 ± 0.8	0.5 ± 0.1	0.440 ± 0.002	2.7	–

1 elements in the average UCC according to [41]

2 elements in soils of the North Caucasus according to [43]

Table 3: The average for 2013-2014 years' elemental content of soils and bottom sediments (BS) for two different stations of Anapa region, upper continental crust (UCC) and average soils of the North Caucasus (SNC) (mean ± standard deviation, µg/g dry weight).

in UCC have the good agreement with the local data for Anapa region (Table 3). For a better description of the local conditions for each sampling site we determined the content of the same elements in soil as well as in sediments. This procedure was used for a more complete analysis of the distribution of elemental content of all considered elements in plants, soil and sediments collected from the Anapa region. Besides the above mentioned statistical analysis techniques, we have also used some graphic analysis procedure such as the ternary diagrams. They allowed to reveal at which extent the content of Cl, Br and I could be used to discriminate the different species of studied plants. All computations were performed using the LibreOffice 5.0.2 and Past 3.0 [42].

Results and Discussion

Accumulation of elements in soils and bottom sediments

The levels of the major and trace elements in soils and sediments from two stations located at 1 km (station 7) and 4 km (station 2c) from the city dump is given in Table 3. The content of the same elements in the average UCC and the levels of some elements in soils of the North Caucasus [43] are listed. Elements in the average UCC according to elements in soils of the North Caucasus according to ref. [43]. The determined concentrations of the majority of elements in soils and BS for each station belonged to close ranges. In that case, we would contemplate these milieus for plants as one. For further analysis, the average values were calculated as arithmetic means for soils (data from surface and from 0-20, 20-40, 40-60 cm layers) and bottom sediments (only from surface). The standard deviation for joint is given on the Figure 3. As follows from Table 3 Cl, Se, Br, and I concentrations in soils and BS from both stations (Figure 2) are higher than in UCC. It can be explained by the location of Anapa region near the sea and the fact that atmospheric supply from the marine environment is the predominant source of these elements in the soil [44,45]. As described in [46] and [47] soil contamination may be considered when concentrations of an element in soils were two- to three times greater than the mean background levels. For our study station 7 (the closest to city dump) hypothetically was the most polluted and the station 2c which situated on the shore was used as background for whole transect. The increasing levels of As, Mo, and Sb in soils and BS from the most polluted station 7 probably indicates the anthropogenic pollution with these elements. Increasing trend of levels of elements from the relatively pristine to polluted area probably ensue from influences of local disposal dump and traffic impacts. The concentrations of all elements (except for V and Ni) reported by for the soils of the North Caucasus are higher than those determined in the soil samples from the most polluted station 7 near city dump of Anapa. Our data were also compared to results of [48] who determined in laboratory conditions the levels of several elements




Figure 3: Average differences between leaves/roots ratios for *Phragmites australis* and *Carex conescens*.

for non-polluted, low polluted and moderate polluted soil from the Southern part of Russia using the integral index of biological state of soil (Table 4). It helps to realize the level of local differences in elemental content of soils from the standard levels for whole region, elements




Figure 4: Studied plant/Reference plant ratios for different parts of *Phragmites australis*, *Carex conescens* and *Cladophora sericea*.

Elements	Soils in Anapa region		Soils in the Southern part of Russia		
	Max	Median	Non-polluted	Low polluted	Moderate polluted
V	150	30	<200	200-300	300-850
Cr	105	30	<70	70-90	90-170
Mn	900	370	<1000	1000-1600	1600-1800
Co	24	4	<18	18-36	36-250
Ni	80	12	<50	50-100	100-700
Zn	270	50	<125	125-200	200-850
As	36.8	7.1	<17	17-30	30-160
Se	2.31	0.25	<0.7	0.7-1.4	1.4-9
Sr	840	510	<250	240-450	450-3200
Mo	15.7	1.1	<8	8-400	>400
Sb	2.1	0.6	<5	5-12	12-200
Ba	690	250	<900	900-1500	1500-4000

Table 4: Maximal and median elemental concentrations ($\mu\text{g/g}$ dry weight) in soils from Anapa region (our data) and values for non-polluted and polluted soils from the Southern part of Russia.

1 elements in soil according to [48]. The concentration of elements which relate to moderate polluted range are given in bold

in soil according to the concentration of elements which relate to moderate polluted range are given in bold. The maximal concentrations of Cr, Zn, As, Se and Sr in soils of Anapa region that were determined at the stations 6 and 7 (the nearest to city dump) are similar with the values reported for moderate polluted soils. Nevertheless, all median values of studied elements Figure 5 in soils of Anapa region are within the range of concentrations determined for non-polluted soils (Table 4) and less than maximum permissible levels of elements established in different countries (Table 5) [49]. Data of maximum permissible levels are widely used for ecological management in assessment of environmental impacts. It was concluded that the soils in study region were in low-polluted state despite the sources of anthropogenic stress.

Accumulation and compartmentalization of elements in water and coastal-aquatic plants

The data about accumulation of elements in different organs of plants were analyzed at the all stations, but after that, the average levels of elemental concentrations for whole Anapa region were calculated as arithmetic mean values obtained from all sampling stations. It helped to realize the ability of different species of plants to reflect the chemical features of environment, including the local pollution influences.

The concentrations of all elements determined (except for K and Cl) are higher in roots of *P. australis* than in leaves. In particular, the leaf/root ratios range from 0.86 for Br to 0.05 for Co. For Sc, V, Fe, Co, I, Cs, Sm and Th the root concentrations are one order of magnitude higher than concentrations in leaf. The obtained results confirm the data that *Phragmites australis* is prevalently a root bioaccumulator species. It is well known that roots are generally the main pathway of trace elements to plants. However, other tissues of *P. australis* in particular, leaves, show the ability readily to translocate such elements as Na, Ti, Zn, Br,

and Sr (Figure 3). In contrast to *P. australis*, the concentration of all elements, except for Fe, Se, Mo, Eu and U are higher in leaves of *Carex conescens* than in roots. Our results emphasized the differences between accumulation features of these two species. The *Carex conescens* could be used as a good bio concentrator of majority of elements from soils and bottom sediments but *P. australis* could be used as a good comparative biomonitor (root type) in clean and polluted areas due to its self-cleaning processes. The obtained results were compared to the available data for *Phragmites*, *Carex*, and *Cladophora*, reported by other authors (Table 6) to represent the variability of concentrations in different regions. The concentrations of most elements in leaves and roots of *Phragmites australis* sampled in the mountain lake in Italy and in the mouth of the longest Sicilian river are higher compared to our results. The exceptions are Ti, Mn, As, Sb and Ti, V, As, Se, which values in roots and leaves, respectively, are higher in the present study. The values of Co, Zn, Rb, and Th in *Carex pendula* sampled in Germany in botanical garden are higher than our data; the reverse trend is observed for Sc, Cr, Fe and Cs. The elemental content of *Cladophora* reported by different authors varies in a wide range depending on the sampling region and the species. Thus, the levels of Mg, Ca and Mn in *Cladophora glomerata* from the lake Karasevov in Siberia are one order of magnitude higher compared to our results [50,51]. In contrast, the content of Ca, Co and Ni in *Cladophora* sp. from the Baltic Sea [52] is one order of magnitude lower than those determined in the present study. The levels of Fe and Zn in *Cladophora glomerata* sampled in the Danube river [53] are 2-fold higher than our data. Thus, the exact concentrations of elements in studied species are absent or not widely available. As a result, it is necessary to determine the range of variability in different pollution conditions. According to the wide variability of elemental content of studied plants across the regions we normalized our data on values for so called reference plant for comparative analysis. The results of normalized elemental concentrations against Reference Plant (RP) show that roots and leaves of *P. australis* are good accumulators of Na, Ti, and Br and, in contrast, contain lower levels of Zn, Rb, and Ba than RP. (Figure 4) In *Carex* roots and leaves the levels of Na, Ti, As, Th, and U are one order of magnitude higher than in RP. In contrast, Mg, K, Mn, Zn, Rb, Cs, and Ba show lower levels in comparison to RP concentrations. The concentrations of the majority of elements in algae




Figure 5: The elemental content of *Phragmites australis*, *Carex conescens* and *Cladophora sericea* normalized against bottom sediments.

Element	Original data			Russia	Germany	Netherlands	USA	Finland
	min	max	median	[48]		[49]		
V	10	150	30	150	-	-	-	100
Cr	6	105	30	90	100	250	1000	100
Mn	150	900	370	1500	-	-	-	-
Co	1.6	24	4	-	50	50	-	20
Ni	3	80	12	85	100	100	-	50
Zn	6	270	50	100	300	500	2500	200
As	3	36.8	7	2	50	30	30	5
Se	0.06	2.31	0.25	-	10	-	-	-
Mo	0.2	15.7	1.1	-	10	40	-	-
Sb	0.1	2.1	0.6	4.5	-	-	-	2
Ba	150	690	250	-	-	400	-	-

Table 5: Maximum permissible levels of elements in soils established in different countries.




Figure 6: Ternary diagram for concentrations of Cl (Cl/10), Br and I for *Phragmites australis*, *Carex conescens* and *Cladophora sericea* normalized against content of these elements in sediments (BS). With some exceptions (*Phragmites* samples), all other points form three distinct clusters corresponding to each type of plant.

Element	<i>Phragmites australis</i>			<i>Carex pendula</i>	<i>Cladophora sp.</i>	<i>Cladophora glomerata</i>	
	Roots (1)	Roots (2,3)	Leaves (2,3)	Whole plant (4)	Whole plant	Whole plant (5)	Whole plant (7)
Na	–	–	–	–	17100	–	3000
Mg	1550	–	–	–	7800	–	23000
Al	–	3153	389	–	–	–	–
K	17000	–	–	–	24500	–	11000
Ca	–	–	–	–	8500	–	170000
Sc	–	–	–	0.19	–	–	–
Ti	–	<0.05	<0.05	–	–	–	–
V	14.5	9.2	<0.05	–	–	–	–
Cr	3.06	6.97	0.69	0.12	–	–	–
Mn	300	475.8	308	–	470	500	18000
Fe	2990	5561	453	460	2400	10000	2300
Co	–	8.0	0.22	0.03	0.5	–	–
Ni	6.52	9.12	1.69	–	3.1	–	6.3
Zn	54	104	28.4	595	60	200	–
As	–	<0.05	<0.05	–	–	–	5
Se	–	<0.5	<0.5	–	–	–	–
Rb	–	–	–	3.8	–	–	–
Sr	–	48.5	82.3	–	90	–	–
Mo	–	16.8	8.5	–	–	–	–
Sb	–	<0.05	<0.05	–	–	–	–
Cs	–	–	–	0.01	–	–	–
Ba	–	47.3	20.1	–	–	–	–
Th	–	–	–	0.45	–	–	–

Table 6: Elemental content of different species of *Phragmites*, *Carex* and *Cladophora* ($\mu\text{g/g}$ dry weight).

Cladophora are at least one order of magnitude higher than in RP. The levels of Zn and Rb, that are lower than RP concentrations, become the exception. (Figure 4) The different composition of *Phragmites australis* and *Carex conescens* with *Cladophora sericea* is explainable by fully different uptake mechanisms of elements either by all surface of plant from water (*Cladophora*) or by roots (*Phragmites* and *Carex*). Also some elements may characterize the different types of plants (for example, algae). Thus, the level of as that is a part of phosphatides in algae and plays an important role in glycometabolism [54] is 140-fold higher in *Cladophora* than its concentration in RP. (Figure 4) The similar patterns of elemental accumulation for all three species are found for several elements. Thus, the levels of Na, Ti and Br are higher than in RP; the reverse trend is revealed for Zn and Rb. It could be explained by abundance or lower concentrations of mentioned elements in the surrounding environment (soils, BS).

Transport of major and trace elements from bottom sediments to plants

The element distributions between the two compartments follow the order: bottom sediment>plant due to differences in concentrations. The differences between species accumulation with taking into account the type of accumulation (roots for *P. australis* and live leaves for *C. conescens*) were represented by normalizing concentrations of elements in plants from the same station on values in bottom sediments. (Figure 5) BS was used as a milieu, which at the same station reflects the local elemental fingerprint of water and other components. It is known that most rooted macrophytes uptake chemicals primarily from sediment pore water but it is also reported that some rooted submersed plants may absorb metals directly from water when they are not readily available in sediments and/or in high concentrations in the surroundings [55]. The one more way of coming the elements to plants is an uptake mechanism of them from air. Plants may absorb Cl, Br, and I directly from the atmosphere; and the marine environment is the main source

of these halogens for plants [45-53]. It is found that the levels of Br and I in algae *Cladophora* are higher than in *Phragmites* and *Carex*. (4) Our results are in agreement with the statement that algae are one of the best accumulators of these elements [56]. To reveal the differences of halogens accumulation in *Phragmites*, *Carex* and *Cladophora* the ternary diagram for the levels of Cl (Cl/10), Br and I in plants normalized against content of these elements in sediments is built. (Figure 6) After that for ternary diagram the values was proportionally reduced to relative units (by using Origin™ 8). *Phragmites* is characterized by high levels of Cl at the majority of sampling sites. In *Carex* the content of Br is equal at all stations except one. *Cladophora* is characterized by high levels of Br and I, while the content of Cl is the minimal. This results demonstrate the specific accumulation features of plants. For example, *Cladophora sericea* accumulates Cl in small relative amounts in comparison to Br and I. *Phragmites australis* in the major cases selects I and Cl regardless Br. In that sense the *Carex conescens* demonstrates the most flexible ability for accumulation of these halogens.

Conclusion

The similarity in elemental concentration in soils and sediments at the majority of sampling stations is established. (Table 3) Sediments act as the primary source of elements for water plants. Regarding Cl, Br and I, the atmospheric supply from the marine environment is the predominant source. The concentration of majority of elements in soils of Anapa region are corresponded to values reported for non-polluted zones. The exception are the most polluted stations (6 and 7) near city dump, where elemental levels are several times higher if compared to median values. The study shows that *Phragmites australis* is prevalently a root bioaccumulator species; in contrast, the concentrations of all elements except for Fe, Se, W, and Mo are higher in leaves of *Carex conescens* than in roots. The different composition of *Cladophora sericea* and *P. australis* with *Carex conescens* is explainable by different elemental uptake, either mainly by entire surface of plant from water (*Cladophora*)

or by roots from sediments (*Phragmites australis* and *Carex conescens*). Translocation of elements varies depending on the physiological property of elemental uptake and is generally more intense through plant tissues than from sediments to plants. Leaves of *P. australis* show the ability to readily translocate such elements as Na, Ti, Zn, Br, and Sr. Among the elements determined the highest translocation between roots and leaves of *Carex conescens* is found for Sc, V, Cr, and Zn. The results of normalized elemental concentrations against Reference Plant show that roots and leaves of *Phragmites australis* are good accumulators of Na, Ti, and Br and, in contrast, contain lower levels of Zn, Rb, and Ba than RP. In *Carex conescens* roots and leaves the levels of Na, Ti, As, Th, and U are one order of magnitude higher than in RP. In contrast, Mg, K, Mn, Zn, Rb, Cs, and Ba show lower levels in comparison to RP concentrations. The concentrations of the majority of elements in algae *Cladophora sericea* are at least one order of magnitude higher than in RP. *Cladophora sericea* accumulated Cl in small relative amounts in comparison to Br and I. *Phragmites australis* in the major cases selected I and Cl regardless Br. In that sense the *Carex conescens* demonstrated the most flexible ability for accumulation of these halogens. The found ratios BS to plants demonstrated the different ability of this three species to reflect the local elemental fingerprints. The levels of majority of elements in *Phragmites australis*, *Carex conescens*, *Cladophora* could be used in future biomonitoring studies on local and regional scales. These plants are potentially useful for monitoring of pollution in general, and for the most elements examined in particular.

References

- Burdin KS (1985) The basis of biological monitoring. Moscow State University Press, p: 160.
- Burdin KS, Gavrilenko EE, Zolotukhina E, Kamnev AN (1988) Study of the macrophytes of the seas of the USSR as biomonitoring objects of heavy metals in the sea water. Ocean Biology, pp: 16-25.
- Burdin KS, Voskoboinikov GM, Zubov EV, Kamnev AN (1993) Marine macro algae as monitoring objects in Arctic ocean bioindication environmental state bioassay and the waste destruction technology. Apatity: MMBI RAS, pp: 72-81.
- Kamnev AN (1989) Structure and functions of the brown algae.
- Vozzinskaya VB, Kamnev AN (1994) Ecological and biological foundations of cultivation and utilization of marine benthic algae.
- Phillips DJH, Rainbow PS (1993) Biomonitoring of trace aquatic contaminants. 2nd edn. Chapman and Hall, London.
- Wang W, Gorsuch JW, Hughes JS (1997) Plants for environmental studies. p: 576.
- Gerhardt A (2000) Biomonitoring of polluted water. Zurich Trans Tech Publ.
- Markert B, Breure T, Zechmeister H (2003) Bioindicators and biomonitors principles concepts and applications. Amsterdam Elsevier.
- Marques JC, Salas F, Patrício J, Teixeira H, Neto JM, et al. (2009) Ecological indicators for coastal and estuarine environmental assessment a user guide. Southampton WIT Press.
- Kamnev AN, Frontasyeva MV, Kravtsova AV, Nekhoroshkov PS (2014) Assessment of elemental composition of macrophytes, soil and bottom sediments along a transect in the coastal zone of Anapa, studied by neutron activation analysis.
- Burdin KS, Zolotuhina E (1998) Heavy Metals in Aquatic Plants (Accumulation and Toxicity). Moscow: Dialog MSU. Russian.
- Kamnev AN, Bunkova OM, Bogatyrev LG (2015) Sabinins ideas and their embodiment. Mineral composition of macrophytes is one of the most important indicator of their contribution to a biogeochemical exchange of the World Ocean. Issues of modern Alology.
- Albers PH, Camardese MB (1993) Effects of acidification on metal accumulation by aquatic plants and invertebrates 1 Constructed wetlands. Environ Toxicol Chem 12: 959-967.
- Bonanno G (2011) Trace element accumulation and distribution in the organs of *Phragmites australis* (common reed) and biomonitoring applications. Ecotoxicology and Environmental Safety 74: 1057-1064.
- Bunkova O, Kamnev A, Yakovlev A, Suhova T (2014) Studying of mineral structure of *Cystoseira barbata* in a sublittoral zone of the northeast coast of the Black Sea. Proceedings of World Conference on Marine Biodiversity Oct 12-16.
- Kravtsova AV, Milchakova NA, Frontasyeva MV (2015) Levels spatial variation and compartmentalization of trace elements in brown algae *Cystoseira* from marine protected areas of Crimea (Black Sea). Marine pollution bulletin 97: 548-554.
- Pankova ES, Kamnev AN, Golubeva EI (2015) Features of the distribution of heavy metals in brown algae *Cystoseira barbata*. International popular science journal Europe-Asia. Earth Sciences 5: 25-28.
- Duman F, Cicek M, Sezen G (2007) Seasonal changes of metal accumulation and distribution in common club rush (*Schoenoplectus lacustris*) and common reed (*Phragmites australis*). Ecotoxicology 16: 457-463.
- Baldantoni D, Ligrone R, Alfani A (2009) Macro- and trace-element concentrations in leaves and roots of *Phragmites australis* in a volcanic lake in Southern Italy. Journal of Geochemical Exploration 101: 166-174.
- Maddison M, Soosaar K, Muring T, Mander U (2009) The biomass and nutrient and heavy metal content of cattails and reeds in wastewater treatment wetlands for the production of construction material in Estonia. Desalination 246: 120-128.
- Bonanno G (2013) Comparative performance of trace element bioaccumulation and biomonitoring in the plant species *Typha domingensis* *Phragmites australis* and *Arundo donax*. Ecotoxicology and Environmental Safety 97: 124-130.
- Strbac S, Sajnovic A, Kasanin GM, Vasic N (2014) Metals in sediment and *Phragmites australis* (common reed) from Tisza river Serbia. Applied Ecology and Environmental Research 12: 105-122.
- Bragato C, Schiavon M, Polese R, Ertani A, Pittarello M, et al. (2009) Seasonal variations of Cu, Zn, Ni and Cr concentration in *Phragmites australis* Trin ex Steud in a constructed wetland of North Italy. Desalination 246: 35-44.
- Ahmad SS, Reshi ZA, Shah MA, Rashid I, Ara R, et al. (2014) Phytoremediation potential of *Phragmites australis* in Hokersar wetland-A Ramsar site of Kashmir Himalaya. International Journal of Phytoremediation 16: 1183-1191.
- Horovitz CT, Schock HH, Horovitz-Kisimova LA (1974) The content of scandium, thorium, silver, and other trace elements in different plant species. Plant and Soil 40: 397-403.
- Pederson JC, Harper KT (1979) Chemical composition of some important plants of southeastern Utah summer ranges related to mule deer reproduction. The Great Basin Naturalist 39: 122-128.
- Ohlson M (1988) Variation in tissue element concentration in mire plants over a range of sites. Ecography 11: 267-279.
- Whitton BA, Burrows IG, Kelly MG (1989) Use of *Cladophora glomerata* to monitor heavy metals in rivers. Journal of Applied Phycology 1: 293-299.
- Malea P, Haritonidis S, Kevrekidis T (1995) Metal content of some green and brown seaweeds from Antikyra Gulf. Hydrobiologia 310: 19-31.
- Chmielewska E, Medved J (2001) Bioaccumulation of heavy metals by green algae *Cladophora glomerata* in a refinery sewage lagoon. Croatica Chemica Acta 74: 135-145.
- Vershinin A, Kamnev A (2001) Harmful algae in Russian European coastal waters. Proceedings of 9th Int Conf on Harmful algal blooms 2000 Feb 7-11 Hobart Australia UNESCO.
- Levkov Z, Krstic S (2003) Use of algae for monitoring of heavy metals in the River Vardar Macedonia. Mediterranean Marine Science 3: 99-102.
- Zbikowski R, Szefer P, Latała A (2007) Comparison of green algae *Cladophora* sp. and *Enteromorpha* sp. as potential biomonitors of chemical elements in the southern Baltic. Science of the total environment 387: 320-332.
- Pogorelov AV, Dulepa SV, Lipilin DA (2013) The space monitoring on the territory of the Krasnodar region. Geomatics 4: 64-71.
- GOST (State Standard) 17.4.4.02-84 Nature Protection. Soils. Methods of Sampling and Preparation of Material for Chemical, Bacteriological, and Helminthological Analysis.
- GOST (State Standard) 17.1.5.01-80 Nature protection. Hydrosphere. General requirements for sampling of bottom sediments of water objects for their

- pollution analysis.
38. Frontasyeva MV (2011) Neutron activation analysis for the Life Sciences. A review *Physics of Elementary Particles and Atomic Nuclei* 42: 332-378.
 39. Dmitriev AY, Pavlov SS (2013) Automated quantitative determination of elements in samples by neutron activation analysis at the IBR-2 at LNP JINR 10: 58-64.
 40. Markert B (1992) Establishing of Reference Plant for inorganic characterization of different plant species by chemical fingerprinting. *Water Air and Soil Pollution*. 64: 533-538.
 41. Rudnick RL, Gao S (2003) Composition of the continental crust. *Treatise on geochemistry* 3: 1-64.
 42. Hammer, Harper DAT, Paul DR (2001) PAST paleontological statistics software package for education and data analysis. *Palaeontologia Electronica* 4: 1-9.
 43. Dyachenko V, Matasova I, Ponomareva O (2014) The trace elements concentrations dynamics in the soil landscapes of the Southern Russia. *Universal Journal of Geoscience* 2: 28-34.
 44. Fuge R (1988) Sources of halogens in the environment, influences on human and animal health. *Environmental Geochemistry and Health* 10: 51-61.
 45. Frontasyeva MV, Steinnes E (2004) Marine gradients of halogens in moss studies by epithermal neutron activation analysis. *Journal of Radioanalytical and Nuclear Chemistry* 261: 101-106.
 46. Logan TJ, Miller RH (1983) Background levels of heavy metals in Ohio farm soils. *Research circular* 275. AGDEX, pp: 508-530.
 47. Chen M, Ma LQ, Harris WG (1999) Baseline concentrations of 15 trace elements in Florida surface soils. *Journal of Environmental Quality* 28: 1173-1181.
 48. Kolesnikov SI, Kazeev KS, Denisova TV, Dadenko EV (2012) Development of a regional ecological regulations of content of contaminants in soils of Southern Russia. *Polythematic Online Scientific Journal of Kuban State Agrarian University* 82: 1-17.
 49. Mynbayeva BN, Imanbekova TG (2013) Assessment of standards of soil's contamination by heavy metals (analytical review). *The Bulletin of the National Academy of Sciences of the Republic of Kazakhstan* 4: 29-39.
 50. Bonanno G, Lo Giudice R (2010) Heavy metal bioaccumulation by the organs of *Phragmites australis* (common reed) and their potential use as contamination indicators. *Ecological Indicators* 10: 639-645.
 51. Maltsev AE, Leonova GA, Bogush AA, Bulycheva TM (2014) Eco geochemical assessment of anthropogenic pollution of flooded open pits ecosystems in Novosibirsk. *Ecology of industrial production* 2: 44-53.
 52. Bojanowski R (1973) The occurrence of major and minor chemical elements in the more common Baltic seaweed. *Oceanologia*. 2: 81-152.
 53. Ravera O (2001) Monitoring of the aquatic environment by species accumulator of pollutants: a review. *Journal of Limnology* 60: 63-78.
 54. Kabata PA, Pendias H (2001) Trace elements in soils and plants. 3rd edn. Boca Raton London New York: CRC Press.
 55. Guilizzoni P (1991) The role of heavy metals and toxic materials in the physiological ecology of submersed macrophytes. *Aquatic Botany*. 41: 87-109.
 56. Saenko GN (1992) Metals and halogens in marine organisms.