

## Role of the COP9 Signalosome in Gastrointestinal Cancers

Sandra Jumpertz<sup>1</sup>, Jürgen Bernhagen<sup>1\*</sup> and Anke K. Schütz<sup>1\*</sup>

<sup>1</sup>Institute of Biochemistry and Molecular Cell Biology, RWTH Aachen University, Pauwelsstraße 30, 52074 Aachen, Germany

\*Corresponding author: Anke K. Schütz, Institute of Biochemistry and Molecular Cell Biology, Pauwelsstraße 30, 52074 Aachen, Germany; Tel: +49 241 80-85096; Fax: +49 241 80-82427; E-mail: [anschuetz@ukaachen.de](mailto:anschuetz@ukaachen.de)

Jürgen Bernhagen, Institute of Biochemistry and Molecular Cell Biology, Pauwelsstraße 30, 52074 Aachen, Germany; Tel: +49 241 80-88831; Fax: +49 241 80-82427; E-mail: [jbernhagen@ukaachen.de](mailto:jbernhagen@ukaachen.de)

Received date: Sep 11, 2014, Accepted date: Dec 27, 2014, Published date: Dec 30, 2014

Copyright: © 2014 Jumpertz S, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

### Abstract

The COP9 signalosome (CSN) is an evolutionarily conserved multi-protein complex found in plants and animals. In mammals, the CSN consists of eight subunits (CSN1-CSN8). It has been suggested to play a key role in tumorigenesis, because its subunits are frequently overexpressed in human cancers and because the CSN is involved in the regulation of a number of processes that are relevant to carcinogenesis and cancer progression, e.g. cell cycle control, signal transduction, and apoptosis. The best-studied biochemical function of the CSN is the control of cellular protein stability via the ubiquitin-proteasome system through regulation of cullin-RING E3 ligase (CRL) activity by deNEDDylation of cullins or by the deubiquitination function of the CSN. Through these activities, the CSN regulates the degradation of several tumor suppressors and oncogenes that are degraded by the 26S proteasome. This review summarizes recent findings that support CSN's role as a potential key player in tumorigenesis in general, but particularly focuses on evidence on the role of the CSN in gastrointestinal cancers. We cover links to tumorigenesis in the liver, stomach, pancreas, and colon, gathering and discussing findings about CSN's expression and its functional impact on cancer development and progression.

**Keywords:** COP9 signalosome; CSN; Gastrointestinal cancer; Hepatocellular carcinoma; Gastric cancer; Pancreatic cancer; Colorectal cancer; JAB1; Deneddylation

### The COP9 Signalosome (CSN)

The constitutive photomorphogenesis 9 (COP9) signalosome (CSN) is an evolutionarily conserved multi-subunit protein complex that regulates pivotal cellular processes such as the cell cycle and apoptosis, as well as a number of oncogenic signal transduction pathways. In 1996, Chamovitz *et al.* first purified the COP9 complex from plants and characterized it as a repressor of dark-grown growth patterns, thus the name “constitutive photomorphogenesis” [1]. An orthologous 450-kDa complex consisting of eight proteins was isolated from human erythrocytes and pig spleen two years later and found to contain the protein c-Jun-activation-domain-binding protein-1 (JAB1) [2,3]. JAB1 had been identified independently of the CSN complex and found to facilitate the binding of c-Jun or JunD to AP-1 sites, thereby acting as a co-activator of AP-1-regulated gene transcription [4]. To date, the CSN has been identified in various eukaryotes, including different fungus species [5-8], *Caenorhabditis elegans* [9], and *Drosophila melanogaster* [10]. The name “COP9 signalosome” and the nomenclature of its subunits was unified for the different species by Deng and colleagues in 2000, designating the subunits CSN1 to CSN8 according to their molecular weight [11]. While monomeric CSN5/JAB1 can exist independently of the complex, the other CSN subunits are exclusively found in the whole complex or as part of smaller sub-complexes [12-14]. CSN subunits 1, 2, 3, 4, 7 and 8 harbor a so-called PCI (proteasome, COP9 signalosome, translation initiation factor) domain whereas CSN5 and CSN6 contain an MPN (Mpr1p-Pad1-N-terminal) domain instead [15,16]. Sharon *et al.* developed a model of the CSN complex based on

a mass spectrometry approach suggesting that the CSN could be organized in two clusters with CSN1/2/3/8 on one side and with CSN4/5/6/7 forming the other cluster. In this model, the two clusters are connected by a single link constituted by a CSN1-CSN6 axis [13]. During the preparation of this article, the crystal structure of the human CSN was resolved at a resolution of 3.8 Å. The structure gives spectacular insight into the architecture of this protein complex and shows that the PCI proteins form an 18-stranded composite  $\beta$ -sheet at the centre of the complex which has a horseshoe-shaped appearance. Located across the PCI ring is a helical bundle of the CSN subunits with the MPN-containing subunits CSN5 and CSN6 at its core, the latter forming an intimate dimer [17]. Notably, only CSN5 contains a JAMM (Jab1/CSN5 MPN metalloenzyme) motif that is placed within its MPN domain, and which is critical for the Zn<sup>2+</sup>-dependent metalloproteinase activity of CSN5 [18]. Through this enzymatic activity, the CSN can cleave NEDD8 (neural precursor cell expressed developmentally down-regulated 8)-conjugates from cullin-RING ligases (CRLs), a process named “deNEDDylation” [19, 20]. CRLs are a large family of multi-subunit E3 ubiquitin ligases assembled on a cullin scaffold. By exchange of various substrate-specific, associated F-box proteins, CRLs have a wide variety of substrates which they ubiquitinate for subsequent proteasomal degradation by the 26S proteasome [21]. By the CSN-dependent removal of NEDD8 from cullins, CRLs are inactivated in short term, as evidenced by biochemical *in vitro* experiments [22,23], whereas *in vivo* experiments in the context of living cells showed that NEDDylation/deNEDDylation cycles are essential for CRL stability and activity [19]. Together, this indicates that the CSN is an important regulator of protein turnover. Moreover, the CSN regulates other E3 ubiquitin ligases such as MDM2 (mouse double minute 2 homolog) and COP1 (constitutively photomorphogenic 1) that target p53, 14-3-3 $\sigma$ , or c-Jun [24-26]. Additionally, the CSN has been shown to associate with

USP15, a deubiquitinase (DUB) involved in NF- $\kappa$ B signaling [27]. Furthermore, by its association with different kinases, the CSN is involved in the phosphorylation of targets such as c-Jun and NF- $\kappa$ B [2]. Recently, the CSN has been linked to diverse disease conditions, e.g. atherosclerosis [28,29], HIV infection [30], cardiomyopathies [31], and microbial sepsis [32].

## The CSN and its Potential Role in Cancer

Cancer diseases still belong to the most challenging human diseases that are associated with high mortality rates and that are difficult to cure with still only limited knowledge about the mechanisms of their development.

The CSN is a potential key player in tumorigenesis and its subunits are often overexpressed in tumors. For example, *CSN5/JAB1* is overexpressed in different tumor entities, e.g. neuroblastoma [33], ovarian tumors [34,35], mamma carcinoma [36,37], hepatocellular carcinoma [38,39], pancreatic cancer [40], lung adenocarcinoma [41], or malignant thyroid lymphoma [42]. The *CSN5* gene is rarely mutated, but its amplification was identified in hepatocellular carcinoma and breast tumors [39,43]. Protein and mRNA analysis in tumor lysates showed that *CSN4* is overexpressed in prostate tumors [44]. An amplification of the *CSN6* gene was identified in human breast cancer samples and in 17 breast cancer cell lines, and CSN6 protein levels were higher in human breast tumors compared to normal breast tissue [45]. In addition, higher CSN6 levels were detected in malignant follicular thyroid carcinomas compared to benign thyroid lesions or normal thyroid tissue [45]. In a transcriptomic analysis, *CSN5* and *CSN6* mRNA levels were shown to be higher in tumors compared to normal tissue for most myelomas as well as in breast cancer and glioblastoma patients [46].

Beyond the altered expression of the CSN and its subunits in tumors, the CSN has an established impact on several important cellular functions that often are dysregulated in tumorigenesis. The CSN is known to be involved in cell cycle and checkpoint control, regulation of proliferation and apoptosis, in the DNA damage repair pathway, and, as discussed above, in the regulation of CRLs and protein degradation. The more general role of the COP9 signalosome in cancer has recently been summarized in a couple of excellent comprehensive reviews. Therefore, we here focus on the CSN and its role in gastrointestinal cancers.

## Molecular Pathways Linking the CSN to Tumorigenesis

Since many key oncogenes and tumor suppressor products such as p27, I $\kappa$ B $\alpha$ , c-Jun, p53, COP1, and 14-3-3 $\sigma$  are degraded via the ubiquitin-26S-proteasome pathway and are directly or indirectly regulated by the CSN [24, 25, 47-49], it is likely that the CSN plays a key role in cancer through several pathways.

CSN5/JAB1 is known to promote the nuclear export and subsequent degradation of the tumor suppressor p53 [50]. p53 is stabilized and activated upon DNA damage, it induces apoptosis in normal cells and is mutated in most tumors, thus incapable of properly regulating cell survival in tumor cells. Furthermore, p53 is a key effector of the DNA damage response in the cell that counteracts DNA damage generated by chemicals, reactive oxygen species, UV or ionizing radiation. Activation of the DNA damage response results in a delay or arrest of the cell cycle and in checkpoint activation and thereby protects the cell against accumulating mutations [51]. Upon UV-induced DNA damage or replication fork stress due to chemicals,

the sensor kinase ATR (ataxia telangiectasia and Rad3-related protein) is activated. ATR subsequently phosphorylates and thereby activates the checkpoint kinase 1 (Chk1). Chk1 is a signal-transducing kinase that has many downstream effectors, e.g. Cdc25, p53, and Cdc45 [52]. The 9-1-1 complex (Rad9-Rad1-Hus1 complex) is loaded onto DNA after genotoxic damage and controls Chk1 phosphorylation. CSN5 was shown to interact with the 9-1-1 complex and to promote its degradation. Thus, CSN5 suppresses checkpoint signaling activation and DNA synthesis recovery after replication stress [53].

The CSN is involved in cell cycle progression at many points. First, CSN5/JAB1 was shown to regulate the CDK-inhibitor p27, promoting its nuclear export and its subsequent degradation – implying a role for the CSN in G1 progression [47]. The E3 ubiquitin ligase SCF<sup>Skp2</sup> controls G1/S cell cycle regulators including p27 [54]. The CSN regulates SCF ligase activity by its deNEDDylase function, but was shown to regulate Skp2 levels directly as well [55]. In addition, Skp2 forms a complex with Cul4A and DDB1 to target p27 [56]. The reduction of CSN levels causes multiple defects in the cell cycle in a subunit- and tissue-specific manner. *Csn5* knockout in T-cells inhibited progression through S phase, but no significant effects on the G1 phase were detected [57]. T-cells with a *Csn8* knockout showed defects in the reentry into the cell cycle from the G0 quiescent stage and the CSN was shown to regulate the transcription of the *G1 cyclins E, D2 and D3*, and of *CDK2* and *CDK4* [58]. Furthermore, a possible direct link between the CSN and G2/M phase progression could be constituted by the interaction of CSN2 with the anaphase-promoting complex (APC)/cyclosome [59]. Additionally, the interaction between MIF and CSN5 was shown to impair the CSN's deNEDDylase function, affecting the activity of SCF ligases involved in G2/M checkpoint control [60]. Of note, the CSN has been suggested to play a key role in cell proliferation and maintenance, because the knockout of any *Csn* subunit tested so far (*Csn2, Csn3, Csn5, Csn6, Csn8*) led to the death of the respective mice at a very early stage during embryonic development due to defects in cell proliferation and survival [45,58,61-63].

Schweitzer *et al.* discovered a functional link between the CSN and NF- $\kappa$ B signaling in the cervix carcinoma cell line HeLa. Here, the CSN has been found to control the activity of CRL <sup>$\beta$ TrCP</sup> that ubiquitinates the NF- $\kappa$ B inhibitor I $\kappa$ B $\alpha$ . Interestingly, the CSN also associated with USP15, thereby promoting I $\kappa$ B $\alpha$  deubiquitination and allowing a precise regulation of NF- $\kappa$ B signaling [27]. A regulative impact of the CSN on NF- $\kappa$ B signaling was also shown in fibroblast cell line HEK293, where the CSN was observed to interact with the I $\kappa$ B $\alpha$  phosphorylating IKK complex and to promote its activity [64]. NF- $\kappa$ B signaling has been mainly associated with inflammatory processes so far, but more and more evidence suggests this signaling pathway to play a key role in tumorigenesis as well and to be an important link of chronic inflammation and cancer [65].

The CSN shows altered expression in different human tumor entities and displays important functions in the cell, implying that the CSN could be a potential target for cancer therapy, although the identified apparent indispensable functions of CSN in cell homeostasis might also contradict this notion. Some of the CSN's functions are mediated by associated kinases as e.g. CK2 $\alpha$ , PKD and IKK2 [64,66]. Interestingly, curcumin effectively inhibits the CSN-associated kinases [67] and is known to be anti-tumorigenic and anti-angiogenic [68,69]. Moreover, especially CSN's regulatory function on the cellular protein degradation machinery could be an attractive target for cancer therapy, as there are already a couple of proteasome inhibitors in

clinical trials and in clinical use. The list encompasses Bortezomib, Carfilzomib, NPI-0052, MLN9708, CEP-18770, ONX0912, and Marizomib. Bortezomib is a peptide boronate inhibitor that binds specifically the catalytic site of the 26S proteasome. It is approved for the treatment of multiple myeloma since several years and its application significantly improved the prognosis of multiple myeloma patients [70,71].

## Role of the CSN in Hepatocellular Carcinoma and in the Liver

Liver cancer is the fifth most common cancer in men and ranks seventh in prevalence for cancers in women. Because of its high fatality (ratio of mortality to incidence) it is the third most common cause of death from cancer worldwide [72]. Hepatocellular carcinoma (HCC) is widespread in Asia and Africa, due to frequent hepatitis B or C infections in these countries. In the Western world, HCC mostly results from liver fibrosis and liver cirrhosis [73]. In an array-based comparative genomic hybridization analysis of HCC patient samples, a copy number gain of chromosome 8q, encoding *CSN5*, was observed in 39% of the cases [39]. In parallel, high *CSN5* expression was detected in these HCC samples compared to normal tissue. By immunohistochemical staining, Patil *et al.* identified 40% of the HCC samples to have *CSN5*-positive nuclei, while they did not detect *CSN5* in normal hepatocytes [39]. Of note, Hsu *et al.* found that *CSN5* overexpression in HCC tissue samples correlates with female gender and hepatitis C infection [38]. Further, *CSN5* expression was monitored in developing HCC; *CSN5* overexpression occurs in early stages of hepatocarcinogenesis and is significantly associated with an expression of MYC-regulated target genes [74]. This could imply that *CSN5* may play a crucial role in liver cancer progression by transcriptional activation of MYC target genes. MYC serves as a transcription factor for a large variety of genes, thereby promoting proliferation, apoptosis, differentiation, metabolism, and genome stability, and is frequently overexpressed in many human cancers [75]. Association of MYC and *CSN5* was described in breast cancer, where *CSN5* promotes transcriptional activity of MYC, but *CSN5* also destabilized MYC via the SCF<sup>Skp2</sup> complex [43]. Besides, a *CSN5* knockdown in HuH7 HCC cells resulted in a significant downregulation of 83 genes that were also significantly elevated in early HCC in comparison to dysplastic nodules (HCC precursors) or cirrhotic (regenerative) nodules [74]. Supporting the hypothesis that *CSN5* overexpression could drive MYC-dependent transactivation and thereby HCC progress, Panattoni *et al.* recently uncovered the relationship of *Csn5* and *Myc* in the regenerating mouse liver [76]. Here, liver-specific *Csn5* deletion leads to replicative stress resulting in cell cycle arrest, polyploidy and apoptosis of hepatocytes. These effects on cell behavior were phenocopied by *c-Myc* overexpression in *Csn5* null hepatocytes of adult mice. Additionally, they determined complex aberrations in cell cycle progression of *Csn5*-deficient hepatocytes: many cells accumulated in the G2/M phase after completion of S phase, but had also a tendency to replicate DNA starting from a hyperdiploid state, resulting in a large fraction of cells with >4N DNA content. These cells underwent massive, uncontrolled DNA re-replication, indicating that the CSN might be a key repressor of DNA re-replication in the course of normal cell proliferation. Together, this study revealed that hepatic *Csn5*, in collaboration with *Myc*, affects the cell cycle and DNA replication [76].

Interestingly, a reduction of the *CSN5* levels by siRNA knockdown resulted in a reduced proliferation or survival rate of many human

HCC cell lines, e.g. Hep3B, HuH7, HepG2, HuH1, and PLC/PRF/5 cells [39,77]. Knockdown of *CSN5* in HuH7 and HepG2 cells was also associated with a strong induction of apoptosis, an accumulation of G0/G1 phase cells and a decrease of cells in S phase, indicating defective cell cycle progression after reduction of *CSN5* in HCC cells. In a gene expression microarray in these HCC cells, Lee *et al.* identified a number of genes affected by *CSN5* knockdown that were also known targets of MYC or of TGFβ1 including *CDK6*, *BRCA1*, and *ATF3*. While genes involved in cell cycle progression, anti-apoptosis, survival and metastasis (*CDK6*, *EIF2S1*, *SLC2A1*, *ITGB1* and *LPL*) were down-regulated after *CSN5* knockdown, the expression of genes associated with pro-apoptotic activity and tumor suppression (*ATF3*, *MPM11*, *GSN*, *TIMP2*, *FSTL3*) was enhanced. Interestingly, depletion of *CSN5* did not affect the protein levels of other CSN subunits, but cullin-1 hyperNEDDylation – dependent on a functional p53 gene – was induced. Moreover, *CSN5* depletion was accompanied by a reduction of the cell cycle regulators cyclin D1, *CDK6*, and *ITGB1*, while the phosphorylation of the TGFβ signaling mediators *SMAD2* and *SMAD3*, which are involved in the regulation of apoptosis, was increased. Furthermore, the knockdown of *CSN5* resulted in a reduction of NF-κB p65, in the downregulation of anti-apoptotic *Bcl-2* and in increased levels of pro-apoptotic *Bak*. Of note, loss of *CSN5* resulted in a decrease of *Skp2* that is part of a CRL and induces the degradation of the *CDK* inhibitors p21 and p27 in HuH7 and HepG2 cells. In parallel, an accumulation of p27 and of the tumor suppressor p53 was observed in HepG2 cells after *CSN5* depletion [77].

To address *CSN5* as a potential target for HCC therapy, *CSN5* siRNA – encapsulated in SNALPs (stable-nucleic-acid-lipid particles) and modified with 2'-O-Methyl or with guanosine nucleosides, was given to mice with HuH7 cell tumors growing in their livers. The mice treated with the siRNA specific for *CSN5* had smaller tumors and an improved well-being compared to control mice, indicating that *CSN5* is an important player in HCC growth and might indeed be a promising target in HCC therapy [77].

Troglitazone is a peroxisome proliferator-activated receptor γ (PPARγ) ligand that exhibits anti-tumoral effects in HCC cells through inhibition of cell proliferation. It leads to an accumulation of p27 and attenuates *Skp2* mRNA expression in HCC cells [78, 79]. Interestingly, treatment of HepG2 cells with troglitazone or another PPARγ ligand, rosiglitazone, downregulates *CSN5* protein levels and mRNA expression in a dose-dependent manner by inhibition of β-catenin/Tcf4 binding to the *CSN5* promoter. As expected, PPARγ inhibitor GW9662 as well as a PPARγ knockdown reversed the troglitazone-induced inhibition of the *CSN5* promoter activity. Inhibition of HepG2 cell growth after troglitazone treatment was partially rescued by ectopic *CSN5* overexpression. Moreover, HepG2 cell xenografts showed a reduced volume and less blood vessels in the tumor microenvironment were detected following troglitazone treatment. In parallel, *CSN5* expression and *CSN5* protein levels were reduced whereas p27 levels were increased in the HepG2-xenografts, after intratumor injection of troglitazone [38].

Topoisomerase II α (topoIIα), an enzyme that loosens up the DNA helix to allow transcription, was found to be overexpressed in HCC and to be associated with chemoresistance [80]. HDAC (histone deacetylase) inhibitors, e.g. AR42 or MS-275, facilitate topoIIα proteolysis through transcriptional activation of the topoIIα-phosphorylating enzyme *CK2α* (casein kinase 2α) by increasing the association of acetylated histone H3 with the *CK2α* promoter [81, 82].



AR42 suppresses HCC tumor growth in mice [83] and enhances complex formation of CSN5 with CK2 $\alpha$  and topoII $\alpha$  in human PLC5 HCC cells [82]. According to this, the knockdown of *CK2a*, of F-box protein *Fbw7* or of *CSN5* reversed the HDAC inhibitor-induced degradation of topoII $\alpha$ , whereas ectopic expression of *CSN5* mimicked the suppressive effects of HDAC-inhibitors on the topoII $\alpha$  levels. Phosphorylation of topoII $\alpha$  by CK2 facilitates its association with the Fbw7-CSN5 complex and its subsequent degradation [82]. Together, these results indicate that high CSN5 levels facilitate topoII $\alpha$  degradation and thus might counteract tumor growth.

Although most CSN/tumor research has focused on CSN5, Yu *et al.* studied CSN3's role in HCC and performed *CSN3* knockdown experiments in HCC cell lines SMMC-7721 and Hep3B. Depletion of *CSN3* resulted in a reduced proliferation rate, inhibition of anchorage-independent growth, cell cycle arrest at G0/G1 phase, inhibited DNA synthesis and enhanced apoptosis of these cells [84] – similar to the results of a *CSN5* knockdown approach in HuH7 and HepG2 cells as mentioned above [77]. Yu *et al.* determined more human HCC samples to be CSN3-positive, especially the nuclei, compared to cirrhotic or normal tissue. In a xenograft model, the knockdown of *CSN3* in SMMC-7721 HCC cells resulted in reduced tumor weight and volume [84]. Interestingly, a *Csn8* knockout in murine hepatocytes induced massive apoptosis of the cells, paralleled by a down-regulation of all other CSN subunits and an impaired deNEDDylation of cullins [85]. As also CSN8 is essential for hepatocyte survival and proliferation, the CSN might represent an interesting target for HCC therapy. Together, these findings indicate that the CSN contributes essentially to HCC tumorigenesis and one might speculate that the COP9 complex as a whole is involved.

Hepatitis is known to be associated with HCC development and HBx (Hepatitis B virus X protein) is a major factor in hepatitis and HCC caused by hepatitis B infections. Interestingly, CSN3, CSN4, and CSN5 were identified as HBx-interacting proteins. In further experiments focusing on CSN5, a direct interaction of HBx and CSN5 was found to facilitate HBx-mediated AP-1 activation in the cervix carcinoma cell line HeLa [86], indicating that the CSN might be a molecular link of hepatitis B virus-induced inflammation of the liver and HCC.

In conclusion, the CSN seems to be overexpressed in HCC, it is involved in maintaining genome stability and controls cell cycle progression, and its function is closely associated with the oncogene MYC, with TGF $\beta$ , and NF- $\kappa$ B signaling in the liver. Of note, HCC cells treated with *CSN5/CSN3*-specific siRNA showed reduced proliferation, enhanced apoptosis, inhibited anchorage-independent growth, and cell cycle arrest, and *in vivo*, siRNA treatment resulted in reduced tumor growth and less angiogenesis. Thus, for future HCC treatment the reduction of CSN levels might be a promising approach.

## Role of the CSN in Gastric Cancer

About one million new cases of stomach cancer were estimated to have occurred in 2008, making it currently the fourth most common malignancy in the world. Gastric cancer is the second leading cause of cancer-related deaths worldwide and its development is very often associated with a *Helicobacter pylori* infection [72,87]. Recently, gene expression profiling of gastric carcinoma patient samples combined with a bioinformatic analysis revealed that CSN5 and CSN6 are significantly upregulated in gastric carcinoma. Furthermore, the transcription factors MYC and MAZ (MYC-associated zinc-finger

protein) showed a similar expression profile as these CSN subunits and were significantly enriched [88].

RUNX3 (Runt-related transcription factor 3) is known to be a strong tumor suppressor in gastric cancer as well as in several other cancer entities. In gastric cancer, the expression of *RUNX3* is frequently lost as a result of promotor hypermethylation, and *RUNX3* is often inactivated by mislocalization in the cell [89,90]. Interestingly, CSN5 and *RUNX3* interact in SNU5 cells, a human gastric cancer cell line, and in HEK293 cells. CSN5's MPN domain and *RUNX3*'s Runt domain were identified to be essential for this interaction. Overexpression of *CSN5* resulted in lower *RUNX3* levels, an effect that is inhibited by the proteasome inhibitor MG132. As expected, a knockdown of *CSN5* led to an increase of *RUNX3* in HEK293 cells. In parallel, CSN5 affected p21 levels – p21 is a target protein of *RUNX3* and a cyclin-dependent kinase inhibitor. In addition, CSN5's MPN domain and its NES region were identified to be essential for the nuclear export and degradation of *RUNX3*. The nuclear export of *RUNX3* is facilitated by CSN5, but is also CRM1 (chromosome region maintenance 1)-dependent. Furthermore, the CSN-associated kinase CK2 $\alpha$  was shown to support the CSN5-mediated degradation of *RUNX3* [91].

MicroRNA-146a (miR-146a) is a modulator of inflammatory signals and was found upregulated in a mouse model of gastric cancer, i.e. the *Gastrin* knockout mouse, and in 2/3 of human gastric adenocarcinomas [92]. In this study, CSN8 and CARD10 (caspase recruitment domain-containing protein 10) were identified as new targets of miR-146a. CSN8, CARD10 and IRAK1 – a known miR-146a target – all belong to signaling pathways leading to NF- $\kappa$ B activation and were all down-regulated after overexpression of miR-146a in SNU638 gastric cancer cells [92]. NF- $\kappa$ B activation is associated with inflammation-mediated gastric carcinogenesis and with gastric cancer cell migration and invasion [93, 94]. The CSN is known to control NF- $\kappa$ B activity by deubiquitination of its inhibitor I $\kappa$ B $\alpha$  – mediated by the CSN-associated deubiquitinase USP15, and by maintaining the activity of I $\kappa$ B $\alpha$  ubiquitin ligase SCF<sup>BTRCP</sup> through its deNEDDylase function [27,29,64,95]. Crone *et al.* suggest the miR-146a levels to be upregulated by NF- $\kappa$ B activity in a negative feedback loop and determined NF- $\kappa$ B activation to be inhibited by overexpression of miRNA-146a in SNU638 cells. Furthermore, CSN2 was found to be down-regulated in this model, indicating that the CSN might be destabilized by miRNA-146a. Notably, the knockdown of *CSN8*, *CSN2* or *CSN5* in SNU638 cells inhibited GPCR-mediated activation of NF- $\kappa$ B comparable to the miR-146a-induced effect [92].

p57 is a CDK inhibitor and its mRNA levels were shown to be very low in gastric cancer cells [96,97]. It shares sequence similarity with p27 and its overexpression causes complete cell cycle arrest in G1 phase [98]. Degradation of p57 is mediated by SCF<sup>Skp2</sup>-dependent ubiquitylation [99]. Of note, recently it was shown that *Skp2* depletion inhibits growth and metastasis of gastric cancer cells by inducing cell cycle arrest in the G1/S phase, paralleled by upregulation of p57 [100]. *CSN6* overexpression resulted in decreased p57 levels and in an accelerated proteasomal degradation of p57 in HEK293 cells. There are hints that CSN6, p57 and Skp2 might form common complexes in the cell, and p57 degradation was shown to be Skp2-dependent [101], indicating that the CSN might associate with SCF<sup>Skp2</sup> E3 ligase to promote p57 ubiquitination and degradation. Thus, one might speculate that CSN(6) promotes the degradation of the CDK inhibitor p57 in gastric cancer cells to promote gastric tumor growth and metastasis.

To date, there are just a few studies addressing CSN's potential role in gastric cancer development, but there are strong hints that the CSN is not just overexpressed in stomach cancer. It rather functionally affects the tumor suppressor RUNX3, the cell cycle regulators p21 and p57 in gastric cancer cells, and it is associated with gastric inflammation by its effect on NF- $\kappa$ B signaling. Altogether, this suggests that the CSN may have an important supportive function in gastric tumorigenesis.

## Role of the CSN in Pancreatic Carcinoma

Pancreatic cancer resulted in about 330,000 deaths worldwide in 2012 and thus is currently the seventh most common cause of deaths due to cancer [102]. Known risk factors for pancreatic cancer, amongst others, are smoking, obesity and diabetes [103, 104]. Pancreatic cancer often has very poor outcomes – 5-year-survival is estimated to be only 5% – due to a lack of symptoms in the early phase of the disease and thereby mostly late diagnosis, when the tumor cells have already metastasized [102].

*CSN5* was shown to be overexpressed in pancreatic adenocarcinoma compared to normal pancreatic tissue [40,105]. In 27 human pancreatic adenocarcinoma cell lines, distinct high-level amplifications of chromosome 7q22 that encodes *CSN6*, were identified [106]. *CSN5* overexpression was found to correlate with absent or low expression of *p27* in pancreatic adenocarcinoma samples, and low *p27* levels are associated with a poor prognosis for patients with pancreatic cancer [40,105]. In most of the tumors, the staining for *CSN5* was predominantly nuclear with a weaker cytoplasmic staining [105]. This inverse correlation of *CSN5* and *p27* can be explained by *CSN5*'s role in promoting nuclear export and subsequent degradation of *p27* [47,107]. In fact, enhancement of proteasomal degradation of *p27* by *CSN5* was seen in different pancreatic cancer cell lines, i.e. Panc-1, Mia PaCa-2, Panc-28. Additionally, a direct interaction between *p27* and *CSN5* was uncovered in a GST-pulldown assay and confirmed in Panc-1 and Panc-28 cells. As expected, a knockdown of *CSN5* in the three mentioned pancreatic cancer cell lines resulted in an accumulation of *p27* and an associated cell cycle arrest. Of note, in Panc-28 cells, it was shown that *CSN5*'s effect on *p27* degradation was Skp2-independent [105]. *p27* degradation is known to be mediated by ubiquitin E3 ligase SCF<sup>Skp2</sup> [108], but recently an alternative E3 ubiquitin ligase, i.e. SIAH-1 (seven in absentia homolog-1), that is regulated by *CSN5*, was shown to interact with *p27* and to regulate its stability [109-111]. Furthermore, *p8* is known to facilitate the translocation of *p27* from the nucleus to the cytoplasm that is mediated by *CSN5*, and *p8* and *CSN5* were shown to directly interact [112]. *p8* is a stress-inducible protein that is known to be overexpressed in stressed pancreatic tissue, in chronic pancreatitis, and in pancreatic cancer [113,114].

SMAD4 is a mediator of TGF $\beta$  signaling, which regulates cell growth and differentiation, normally acts as a tumor suppressor, and was shown to be mutated or even inactivated in half of pancreatic cancers [115]. SMAD4 protein levels appear low in human pancreatic ductal adenocarcinoma [116], and SMAD4 inactivation has been suggested to be a key event during pancreatic tumorigenesis [117]. Remarkably, *CSN5* promotes SMAD4 degradation in pancreatic cancer cell lines Panc-1 and AsPC-1 via SCF <sup>$\beta$ TrCP</sup> E3 ubiquitin ligase [116,118]. Point-mutated SMAD4, typically found in pancreatic adenocarcinoma, showed an increased affinity to the F-box protein  $\beta$ TrCP; thus its degradation that is additionally promoted by high *CSN5* levels, is potentially enhanced in pancreatic tissue [116].

As already discussed for HCC, the interplay between *MYC* and *CSN5* might play a key role in pancreatic cancer as well. In Mia PaCa-2 and Panc-1 cells, knockdown of *CSN5/JAB1* impaired cell proliferation and enhanced apoptosis – independent of the *p53* status of the cells. These *CSN5*-depleted cells showed increased *MYC* levels and an additional depletion of *MYC* rescued cells from growth suppression [119].

Interestingly, Li *et al.* tested PEGylated curcumin, an inhibitor of the CSN-associated kinases, and asked if it inhibited cell growth of pancreatic cancer cells, and thus if it might be a potential future therapeutic for pancreatic cancer. PEG (polyethylene glycol)-ylation makes curcumin, that is known to be anti-tumorigenic, water-soluble and thereby facilitates its delivery. In this study, (PEG-)curcumin inhibited cell growth of pancreatic Panc-1, Mia PaCa-2, BxPC-3, and AsPC-1 cells in a dose-dependent manner. Additionally, the CRL targets *p27* and SMAD4 were upregulated after (PEG-)curcumin treatment in Panc-1 and AsPC-1 cells, indicating that *CSN*'s activity is inhibited. Of note, PEG-curcumin sensitized pancreatic cancer cells to gemcitabine-induced apoptosis and inhibition of cell proliferation [120]; this supports the hypothesis that the CSN and its associated kinases are promising targets in future pancreatic cancer therapy.

Together, in pancreatic tissue the CSN regulates the cell cycle by its close association with *p27*; it also affects TGF $\beta$  signaling and the tumor suppressor SMAD4, and functionally interacts with the oncogene *MYC*. As shown by Li *et al.*, an inhibition of *CSN* activity or its associated kinases might have a therapeutic benefit for pancreatic cancer patients.

## Role of the CSN in Colorectal Carcinoma

Colorectal cancer (CRC) is the third most common cancer in men and the second most prevalent one in women worldwide, resulting in about 608,000 deaths related to CRC in 2008 [72]. CRC is a very common form of cancer in the Western world due to lifestyle habits as smoking, excessive alcohol use or obesity, which are known risk factors. Diagnosis is often delayed; consequently, CRC is one of the deadliest forms of cancer.

To date, there are just a few studies that have addressed the expression and the potential role of the CSN in colorectal carcinoma. A recent study showed that 70% of the patients with colon adenocarcinoma have significantly higher *CSN5* mRNA levels in the tumor tissue compared to surrounding normal tissue, and that 40% of them have significantly higher *CSN6* mRNA levels in their tumors [46]. Next to this expression study, there are first hints from *in vitro* studies that the CSN has a functional impact on CRC cell behavior and that it might contribute to the development of the 'hallmarks' of this cancer entity. Schütz *et al.* showed that the reduction of *CSN5* levels correlated with a reduction of *CSN1* and *CSN8* protein levels in SW480 CRC cells, and that the proliferation rate of different CRC cell lines (SW480, HCT116, HT29, CaCo2) was reduced after *CSN5* knockdown. Surprisingly, also the rate of apoptosis after etoposide treatment was reduced by *CSN5* knockdown in HCT116 cells [121]. During CRC tumorigenesis, mutations or deletions of a number of tumor suppressor genes and oncogenes accumulate [122]. CRC development is often associated with mutations of the Wnt signaling pathway. Mostly, the *APC* gene is mutated, but mutations in  $\beta$ -catenin/*CTNBN1* are also observed in CRC [123, 124].

In normal cells,  $\beta$ -catenin levels are tightly regulated by a cytoplasmic destruction complex mainly consisting of APC, Axin, and


GSK3 $\beta$ , which phosphorylates  $\beta$ -catenin, marking it for ubiquitination and degradation by the 26S proteasome. When Wnt signaling is activated by binding of a Wnt ligand to a receptor of the Frizzled family, degradation of  $\beta$ -catenin is inhibited,  $\beta$ -catenin accumulates in the cytoplasm and enters the nucleus to promote transcription of several target genes together with TCF/LEF transcription factors [125]. Activated Wnt signaling is known to be pro-proliferative and several Wnt target genes are known to promote tumorigenesis, e.g. MYC, COX-2, CD44, survivin or cyclin D1. When APC is mutated in intestinal cells and thus becomes dysfunctional,  $\beta$ -catenin degradation is impaired and Wnt signaling is permanently 'on', resulting, amongst other effects, in enhanced cell proliferation. Alternatively, e.g. in HCT116 cells,  $\beta$ -catenin itself can be mutated and its degradation becomes ineffective [124].

The CSN was shown to affect the protein levels of  $\beta$ -catenin in CRC cells. After knockdown of *CSN5*, (phospho-) $\beta$ -catenin levels were reduced in HeLa and SW480 cells, indicating that the CSN might somehow 'stabilize'  $\beta$ -catenin. However interestingly, in SW480 cells, CSN5 was shown to also promote short-term  $\beta$ -catenin degradation in a cycloheximide assay [121]. Similar results were obtained with a knockdown of *CSN1* that resulted in a retarded degradation of  $\beta$ -catenin in HeLa cells [126]. Additionally, a knockdown of *CSN5* resulted in reduced Wnt/ $\beta$ -catenin target gene expression in HCT116 cells, as shown for *Axin2*, *MYC*, *CD44* and *BMP4*, indicating that indeed, functional Wnt signaling is affected [110]. In line with this notion, Xin *et al.* observed reduced COX-2 protein levels in HT29 cells after knockdown of *CSN5* [127]. COX-2 is a Wnt target gene that is upregulated in CRC and contributes to its formation. It induces angiogenesis, inhibits apoptosis and promotes tumor cell invasiveness. Of note, COX-2 inhibitors are already used for CRC prevention in FAP (familial adenomatous polyposis) patients [128,129].

The effect of CSN5 on  $\beta$ -catenin could be an effect on the canonical Wnt pathway and associated  $\beta$ -catenin degradation. For example, the CSN, through its deNEDDylase function, might regulate the activity of the E3 ligase CRL $^{\beta$ TrCP, which is responsible for the ubiquitination of  $\beta$ -catenin. Alternatively, it was shown that a knockdown of *CSN1* in HeLa cells resulted in an accelerated degradation of APC, indicating that the canonical degradation of  $\beta$ -catenin might be impaired due to the reduced APC levels following reduction of the CSN. Furthermore, CSN-associated USP15 stabilized APC and thus might destabilize  $\beta$ -catenin in HeLa cells [126]. Huang *et al.* performed density-gradient centrifugation experiments and found that CSN3,  $\beta$ -catenin, and cullin-1 co-sedimented in the same fractions. By FLAG-pull-downs, they also precipitated CSN2 together with components of the  $\beta$ -catenin destruction complex and of CRL $^{\beta$ TrCP and proposed the formation of a  $\beta$ -catenin-degrading super-complex containing the 26S proteasome, CRL $^{\beta$ TrCP, the canonical  $\beta$ -catenin destruction complex, and the CSN [126].

On the other hand, it was recently shown that CSN5 affects SIAH-1, a key mediator of an alternative  $\beta$ -catenin degradation pathway [110]. SIAH-1 is induced by p53 and forms an alternative degradation complex for  $\beta$ -catenin together with SIP, SKP1, and EBI [130,131], and was shown to promote  $\beta$ -catenin degradation in HCT116 and other cells [110,131]. Of note, in HCT116 CRC cells, CSN5 was shown to affect SIAH-1 protein levels, as after *CSN5* knockdown SIAH-1 levels were reduced and, vice versa after *CSN5* overexpression, SIAH-1 levels were elevated. CSN5 promoted *SIAH-1* expression and SIAH-1 protein degradation – presumably by its deNEDDylase activity. SIAH-1,  $\beta$ -catenin and CSN5 potentially form common protein

complexes and thus might also functionally interact as indicated by *SIAH-1/CSN5*-double-knockdown experiments with an analysis of Wnt target gene expression [110]. Recently, Tanaka *et al.* determined Sec6, a component of the exocyst complex, to interact with SIAH-1 and CSN5 in HEK293 cells. Sec6 was shown to regulate the cytoplasmic translocation of p27 and to promote its degradation [109]. Thus, the interplay between CSN(5) and SIAH-1 might be essential for the regulation of  $\beta$ -catenin and p27 levels, and thereby might affect cell cycle regulation and proliferation of cancer cells.



**Figure 1:** Proteins and their associated cellular processes affected by the COP9 signalosome in gastrointestinal cancers. Proteins depicted as circles are arranged according to their related cellular processes, divided into six categories: cell cycle, angiogenesis, inflammation, proliferative signaling, apoptosis and invasion, which are reflecting the hallmarks of cancer to a large part. Proteins located at the border of two categories were found to play a role in both cellular processes. For reasons of clarity, COX-2, NF- $\kappa$ B, and p53, which are related to multiple processes, are depicted twice. Abbreviations: CycD1 (Cyclin D1), topoIIa (topoisomerase IIa), COX-2 (cyclooxygenase-2), NF- $\kappa$ B (nuclear factor  $\kappa$ B), MYC (v-myc avian myelocytomatosis viral oncogene homolog), SIAH-1 (seven in absentia homolog-1), SMAD2/3/4 (small mothers against decapentaplegic 2/3/4), RUNX3 (Runt-related transcription factor 3).

Hyperthermic intraperitoneal chemotherapy (HIPEC) in combination with complete cytoreductive surgery provides the only chance for long-term survival for selected patients with a variety of peritoneal neoplasms with digestive malignancies [132]. In HIPEC therapy, mitomycin C often is used and functions to induce DNA damage and apoptosis in tumor cells. The CSN positively influences the DNA damage response pathway that is induced by mitomycin C and counteracts chemotherapy. In a HIPEC model with HT29 cells, mitomycin C induced elevated levels of CSN1, CSN3, CSN5, and CSN8, and accelerated the deNEDDylation of cullin-1. When HT29 cells were treated with curcumin, an unspecific inhibitor of the CSN-



associated kinases, the mitomycin C-induced apoptosis rate increased. In parallel, a mimic of let-7a-1, a micro-RNA, was tested to support the chemotherapeutic effects of mitomycin C. The let-7a-1 mimic treatment resulted in a reduced expression of different CSN subunits in HT29 and HeLa cells and, as a potential consequence, led to increased levels of the CDK inhibitor p27 and of the tumor suppressor p53 in HT29 cells [133,134]. In contrast, knockdown of *CSN5* resulted in a small but significant decrease of p53 levels in SW480 cells [121].

Loda *et al.* correlated a lack of p27 in colorectal carcinoma to a poor survival prognosis compared to CRC patients whose tumors expressed p27 [135]. For breast and pancreatic cancer, it was shown that the expression of *CSN5* negatively correlated with the levels of p27 [36, 37, 105]. According to these data, one would expect high *CSN5* levels in CRC tumors to correlate with low p27 levels and with a poor prognosis. However, in a recent study no significant effect of *CSN5* on p27 protein levels in SW480 CRC cells was observed [121], and to our knowledge there is no study indicating the otherwise established p27-*CSN5* correlation would be relevant for CRC development.

In summary, the CSN and in particular its subunit *CSN5* affects many key effectors and signaling pathways in cancer cells. Some of these effects as outlined in this review article are presumably relevant in a more general sense, whereas others appear to be more specific for a certain tissue, cell type, or tumor entity.

In colorectal cancer, the CSN might be a key player as well driving tumorigenesis, because it affects signaling pathways that are crucial for CRC development, such as the Wnt signaling pathway, and is involved in  $\beta$ -catenin degradation. Similarly, *CSN5* depletion results in reduced proliferation of CRC cells, and, as the CSN is an important player in the DNA damage response, it might be a promising therapeutic target

for CRC therapy, e.g. in combination with mitomycin C chemotherapy. On the other hand, it should be kept in mind that evidence linking the CSN to CRC has mainly been *in vitro* so far.

## Conclusion

A number of studies over the last several years have established a likely functional role of the COP9 signalosome in several gastrointestinal cancers, i.e. HCC, pancreatic cancer, gastric cancer, and CRC (summarized in table 1). There is strong evidence that the CSN plays a key role in regulating the cell cycle, proliferation, and apoptosis, and there are hints that the CSN also affects additional key processes driving tumorigenesis including angiogenesis, invasion, and chronic inflammation (Figure 1). Whether the CSN also supports the acquirement of further hallmarks of cancer, e.g. metastasis, replicative immortality, or deregulation of cellular energetics, needs to be elucidated. As outlined in this review article, the CSN affects different cellular signaling pathways such as Wnt, TGF $\beta$ , and NF- $\kappa$ B signaling and regulates the protein levels of important tumor suppressors. Overall, the described functions of the CSN seem to be in line with a pro-tumorigenic phenotype, as e.g. shown in many tumors after overexpression of different CSN subunits. Moreover, in preclinical *in vivo* models, a reduction of the CSN resulted in reduced tumor volume and less blood vessels, pointing to a potential therapeutic benefit of agents that reduce CSN protein levels and/or inhibit its functions in tumors. Thus, the CSN might be an interesting target for future therapeutic approaches for the treatment of cancer, while care needs to be taken to not interfere with its inherently crucial functions in cellular physiology.

Organ / tumor entity	CSN subunit	Cancer (cell) type	Effect	Reference
Liver	CSN3	SMMC-7721 and Hep3B cells	CSN3 promotes proliferation, anchorage-independent growth, cell cycle and DNA-synthesis and counteracts apoptosis	[84]
		Xenograft mouse model (SMMC-7721 cells)	CSN3 knockdown reduces tumor growth and weight	[84]
		Human HCC tissue samples	CSN3 is overexpressed and accumulates in the nucleus	[84]
	CSN5	Hep3B, HuH7, HepG2, HuH1 and PLC/PRF/5 cells	CSN5 promotes cell proliferation	[39,77]
		HuH7 and HepG2 cells	CSN5 facilitates cell cycle and counteracts apoptosis, decreases phosphorylation of SMAD2 and SMAD3 (key mediators of TGF $\beta$ signaling) and enhances NF- $\kappa$ B p65 levels	[77]
		HuH7 cells	CSN5 siRNA encapsulated in SNALP impairs survival	[77]
		HuH7 cells	CSN5 knockdown downregulates 83 genes shown to be elevated in early HCC	[74]
		HepG2 cells	PPAR $\gamma$ ligands downregulate CSN5 protein levels and mRNA expression	[38]
		PLC5 cells	CSN5 promotes the HDAC inhibitor-induced degradation of topol $\alpha$ by CRL <sup>Fbw7</sup>	[82]
		Xenograft mouse model (HepG2 cells)	PPAR $\gamma$ ligand treated tumors are smaller and have less blood vessels, correlating with increased expression of <i>CSN5</i> and p27	[38]
Xenograft mouse model (HuH7 cells)	CSN5 siRNA encapsulated in SNALP decreases tumor growth and improves well-being of the animals	[77]		

		Human HCC tissue samples	Chromosome 8q, encoding CSN5, gains copy number in 39% of samples, nuclear CSN5 levels are enhanced.	[39]
			CSN5 overexpression correlates with female gender and hepatitis C viral infection	[38]
			CSN5 overexpression occurs in early stages of hepatocarcinogenesis and is associated with expression of MYC-regulated target genes	[74]
Stomach	CSN2	SNU638 cells	CSN2 knockdown inhibits GPCR-mediated NF-κB activation	[92]
	CSN5	SNU638 cells	CSN5 knockdown inhibits GPCR-mediated NF-κB activation	[92]
		SNU5 cells	CSN5 interacts with RUNX3	[136]
		Human gastric carcinoma tissue samples	Enhanced levels of CSN5 correlate with CSN6 levels and MYC and MAZ expression	[88]
	CSN6	Human gastric carcinoma tissue samples	Enhanced levels of CSN6 correlate with CSN5 levels and MYC and MAZ expression	[88]
CSN8	SNU638 cells	Overexpression of miRNA-146a downregulates CSN8 and CARD10	[92]	
Pancreas	CSN5	Mia PaCa-2, Panc-28 and Panc-1 cells	CSN5 facilitates p27 degradation	[105]
		Panc-28 and Panc-1 cells	CSN5 interacts with p27	[105]
		Mia PaCa-2 and Panc-28 cells	CSN5 promotes cell cycle progression	[105]
		Panc-28 cells	CSN5-dependent degradation of p27 might be Skp2-independent	[105]
		Panc-1 and AsPC-1 cells	CSN5 promotes SMAD4 degradation via SCF <sup>βTrCP</sup>	[116,118]
		Mia PaCa-2 and Panc-1 cells	CSN5 promotes cell proliferation and inhibits apoptosis via MYC	[119]
		Panc-1 and AsPC-1 cells	PEGylated curcumin (CSN-associated kinase inhibitor) promotes apoptosis following gemcitabine treatment, inhibits cell growth via decelerating cell proliferation and increases levels of CSN5 degradation targets p27 and SMAD4	[120]
		Mia PaCa-2 cells	PEGylated curcumin inhibits cell growth via decelerating proliferation	[120]
		BxPC-3 cells	PEGylated curcumin inhibits cell growth	[120]
		Pancreatic adenocarcinoma tissue samples	CSN5 is overexpressed in tumors, mainly in the cytoplasm; patients with p27-positive tumors had a significant better prognosis than those with p27-negative tumors, and p27 localized exclusively in the nucleus further improved their prognosis	[40]
	Pancreatic adenocarcinoma tissue samples	CSN5 is overexpressed and inversely correlates with p27 levels; in 72% of the tumors predominant nuclear staining for CSN5; tumors with predominant cytoplasmic CSN5-staining were all p27-negative and had a higher proliferation index than cells with predominant nuclear CSN5-staining	[105]	
CSN6	Pancreatic adenocarcinoma tissue samples	High-level amplifications of chromosome 7q22, encoding CSN6	[106]	
Colon	CSN5	SW480 cells	CSN5 knockdown reduces (phospho-)β-catenin, CSN1, CSN8, and p53 protein levels, and decelerates proliferation	[121]
		HCT116 cells	CSN5 promotes apoptosis after etoposide treatment and enhances proliferation	[121]
		HCT116 cells	CSN5 promotes expression of Wnt/β-catenin target genes and of <i>SIAH-1</i> , but also supports <i>SIAH-1</i> degradation, depending on Cullin1-NEDDylation	[110]
		HT29 cells	CSN5 knockdown affects COX-2 protein levels	[127]



		HT29 cells	CSN5 promotes proliferation	[121]
		CaCo2 cells	CSN5 promotes proliferation	[121]
		Colon adenocarcinoma tissue samples	Higher CSN5 mRNA levels in 70% of the tumor samples compared to normal tissue	[46]
	CSN6	Colon adenocarcinoma tissue samples	Higher CSN6 mRNA levels in 40% of the tumor samples compared to normal tissue	[46]

**Table 1:** Summary of functional evidence for a role of the CSN in gastrointestinal tumorigenesis.

## Acknowledgements

We are indebted to numerous colleagues for inspiring and helpful discussions and apologize to the many colleagues and scientists in the field whose work we were not able to cite due to space constraints.

## Financial support

This work was funded by a START-Up grant of the DFG excellence program of RWTH Aachen University (grant 351400) and a START grant of RWTH Aachen University Medical School (grant 690644) to A.K.S., by the START consortium grant 69605-5/project P6 of RWTH Aachen University Medical School to J.B., and by the IZKF Aachen grant SP/O-3 to J.B..

## References:

- Chamovitz DA, Wei N, Osterlund MT, von Arnim AG, Staub JM, et al. (1996) The COP9 complex, a novel multisubunit nuclear regulator involved in light control of a plant developmental switch. *Cell* 86: 115-121.
- Seeger M, Kraft R, Ferrell K, Bech-Otschir D, Dumdey R, et al. (1998) A novel protein complex involved in signal transduction possessing similarities to 26S proteasome subunits. *FASEB J* 12: 469-78.
- Wei N, Deng XW (1998) Characterization and purification of the mammalian COP9 complex, a conserved nuclear regulator initially identified as a repressor of photomorphogenesis in higher plants. *Photochem Photobiol* 68: 237-241.
- Claret FX, Hibi M, Dhut S, Toda T, Karin M (1996) A new group of conserved coactivators that increase the specificity of AP-1 transcription factors. *Nature* 383: 453-457.
- Busch S, Eckert SE, Krappmann S, Braus GH (2003) The COP9 signalosome is an essential regulator of development in the filamentous fungus *Aspergillus nidulans*. *Mol Microbiol* 49: 717-730.
- Maytal-Kivity V, Pick E, Piran R, Hofmann K, Glickman MH (2003) The COP9 signalosome-like complex in *S. cerevisiae* and links to other PCI complexes. *Int J Biochem Cell Biol* 35: 706-715.
- Mundt KE, Porte J, Murray JM, Brikos C, Christensen PU, et al. (1999) The COP9/signalosome complex is conserved in fission yeast and has a role in S phase. *Curr Biol* 9: 1427-1430.
- Wee S, Hetfeld B, Dubiel W, Wolf DA (2002) Conservation of the COP9/signalosome in budding yeast. *BMC Genet* 3: 15.
- Luke-Glaser S, Roy M, Larsen B, Le Bihan T, Metalnikov P, et al. (2007) CIF-1, a shared subunit of the COP9/signalosome and eukaryotic initiation factor 3 complexes, regulates MEL-26 levels in the *Caenorhabditis elegans* embryo. *Mol Cell Biol* 27: 4526-4540.
- Freilich S, Oron E, Kapp Y, Nevo-Caspi Y, Orgad S, et al. (1999) The COP9 signalosome is essential for development of *Drosophila melanogaster*. *Curr Biol* 9: 1187-1190.
- Deng XW, Dubiel W, Wei N, Hofmann K, Mundt K, et al. (2000) Unified nomenclature for the COP9 signalosome and its subunits: an essential regulator of development. *Trends Genet* 16: 202-203.
- Kato JY, Yoneda-Kato N (2009) Mammalian COP9 signalosome. *Genes Cells* 14: 1209-1225.
- Sharon M, Mao H, Boeri Erba E, Stephens E, Zheng N, et al. (2009) Symmetrical modularity of the COP9 signalosome complex suggests its multifunctionality. *Structure* 17: 31-40.
- Fukumoto A, Tomoda K, Kubota M, Kato JY, Yoneda-Kato N (2005) Small Jab1-containing subcomplex is regulated in an anchorage- and cell cycle-dependent manner, which is abrogated by ras transformation. *FEBS Lett* 579: 1047-1054.
- Hofmann K, Bucher P (1998) The PCI domain: a common theme in three multiprotein complexes. *Trends Biochem Sci* 23: 204-205.
- Wei N, Serino G, Deng XW (2008) The COP9 signalosome: more than a protease. *Trends Biochem Sci* 33: 592-600.
- Lingaraju GM, Bunker RD, Cavadini S, Hess D, Hassiopen U, et al. (2014) Crystal structure of the human COP9 signalosome. *Nature* 512: 161-165.
- Ambroggio XI, Rees DC, Deshaies RJ (2004) JAMM: a metalloprotease-like zinc site in the proteasome and signalosome. *PLoS Biol* 2: E2.
- Cope GA, Suh GS, Aravind L, Schwarz SE, Zipursky SL, et al. (2002) Role of predicted metalloprotease motif of Jab1/Csn5 in cleavage of Nedd8 from Cul1. *Science* 298: 608-611.
- Lyapina S, Cope G, Shevchenko A, Serino G, Tsuge T, et al. (2001) Promotion of NEDD-CUL1 conjugate cleavage by COP9 signalosome. *Science* 292: 1382-1385.
- Petroski MD, Deshaies RJ (2005) Function and regulation of cullin-RING ubiquitin ligases. *Nat Rev Mol Cell Biol* 6: 9-20.
- Saha A, Deshaies RJ (2008) Multimodal activation of the ubiquitin ligase SCF by Nedd8 conjugation. *Mol Cell* 32: 21-31.
- Read MA, Brownell JE, Gladysheva TB, Hottelet M, Parent LA, et al. (2000) Nedd8 modification of cul-1 activates SCF(beta(TrCP))-dependent ubiquitination of IkkappaBalpha. *Mol Cell Biol* 20: 2326-2333.
- Zhang XC, Chen J, Su CH, Yang HY, Lee MH (2008) Roles for CSN5 in control of p53/MDM2 activities. *J Cell Biochem* 103: 1219-1230.
- Choi HH, Gully C, Su CH, Velazquez-Torres G, Chou PC, et al. (2011) COP9 signalosome subunit 6 stabilizes COP1, which functions as an E3 ubiquitin ligase for 14-3-3sigma. *Oncogene* 30: 4791-4801.
- Bianchi E, Denti S, Catena R, Rossetti G, Polo S, et al. (2003) Characterization of human constitutive photomorphogenesis protein 1, a RING finger ubiquitin ligase that interacts with Jun transcription factors and modulates their transcriptional activity. *J Biol Chem* 278: 19682-19690.
- Schweitzer K, Bozko PM, Dubiel W, Naumann M (2007) CSN controls NF-kappaB by deubiquitinylation of IkkappaBalpha. *EMBO J* 26: 1532-1541.
- Azuma Y, Takada M, Maeda M, Kioka N, Ueda K (2009) The COP9 signalosome controls ubiquitinylation of ABCA1. *Biochem Biophys Res Commun* 382: 145-148.
- Asare Y, Shagdarsuren E, Schmid JA, Tilstam PV, Grommes J, et al. (2013) Endothelial CSN5 impairs NF-kB activation and monocyte adhesion to endothelial cells and is highly expressed in human atherosclerotic lesions. *Thromb Haemost* 110: 141-152.

30. Kinoshita SM, Krutzik PO, Nolan GP (2012) COP9 signalosome component JAB1/CSN5 is necessary for T cell signaling through LFA-1 and HIV-1 replication. *PLoS One* 7: e41725.
31. Su H, Li J, Menon S, Liu J, Kumarapeli AR, et al. (2011) Perturbation of cullin deneddylation via conditional Csn8 ablation impairs the ubiquitin-proteasome system and causes cardiomyocyte necrosis and dilated cardiomyopathy in mice. *Circ Res* 108: 40-50.
32. Deng Z, Pardi R, Cheadle W, Xiang X, Zhang S, et al. (2011) Plant homologue constitutive photomorphogenesis 9 (COP9) signalosome subunit CSN5 regulates innate immune responses in macrophages. *Blood* 117: 4796-4804.
33. Shen L, Tsuchida R, Miyauchi J, Saeki M, Honna T, et al. (2000) Differentiation-associated expression and intracellular localization of cyclin-dependent kinase inhibitor p27KIP1 and c-Jun co-activator JAB1 in neuroblastoma. *Int J Oncol* 17: 749-54.
34. Sui L, Dong Y, Ohno M, Watanabe Y, Sugimoto K, et al. (2001) Jab1 expression is associated with inverse expression of p27(kip1) and poor prognosis in epithelial ovarian tumors. *Clin Cancer Res* 7: 4130-4135.
35. Sui L, Dong Y, Watanabe Y, Yamaguchi F, Sugimoto K, et al. (2006) Clinical significance of Skp2 expression, alone and combined with Jab1 and p27 in epithelial ovarian tumors. *Oncol Rep* 15: 765-771.
36. Esteva FJ, Sahin AA, Rassidakis GZ, Yuan LX, Smith TL, et al. (2003) Jun activation domain binding protein 1 expression is associated with low p27(Kip1) levels in node-negative breast cancer. *Clin Cancer Res* 9: 5652-5659.
37. Kouvaraki MA, Rassidakis GZ, Tian L, Kumar R, Kittas C, et al. (2003) Jun activation domain-binding protein 1 expression in breast cancer inversely correlates with the cell cycle inhibitor p27(Kip1). *Cancer Res* 63: 2977-2981.
38. Hsu MC, Huang CC, Chang HC, Hu TH, Hung WC (2008) Overexpression of Jab1 in hepatocellular carcinoma and its inhibition by peroxisome proliferator-activated receptor[gamma] ligands in vitro and in vivo. *Clin Cancer Res* 14: 4045-4052.
39. Patil MA, Gutgemann I, Zhang J, Ho C, Cheung ST, et al. (2005) Array-based comparative genomic hybridization reveals recurrent chromosomal aberrations and Jab1 as a potential target for 8q gain in hepatocellular carcinoma. *Carcinogenesis* 26: 2050-2057.
40. Fukumoto A, Ikeda N, Sho M, Tomoda K, Kanehiro H, et al. (2004) Prognostic significance of localized p27Kip1 and potential role of Jab1/CSN5 in pancreatic cancer. *Oncol Rep* 11: 277-284.
41. Goto A, Niki T, Moriyama S, Funata N, Moriyama H, et al. (2004) Immunohistochemical study of Skp2 and Jab1, two key molecules in the degradation of P27, in lung adenocarcinoma. *Pathol Int* 54: 675-681.
42. Ito Y, Yoshida H, Matsuzuka F, Matsuura N, Nakamura Y, et al. (2003) Jun activation domain-binding protein 1 expression in malignant lymphoma of the thyroid: its linkage to degree of malignancy and p27 expression. *Anticancer Res* 23: 4121-4125.
43. Adler AS, Lin M, Horlings H, Nuyten DS, van de Vijver MJ, et al. (2006) Genetic regulators of large-scale transcriptional signatures in cancer. *Nat Genet* 38: 421-430.
44. Bhansali M, Shemshedini L (2014) COP9 subunits 4 and 5 target soluble guanylyl cyclase  $\alpha 1$  and p53 in prostate cancer cells. *Mol Endocrinol* 28: 834-845.
45. Zhao R, Yeung SC, Chen J, Iwakuma T, Su CH, et al. (2011) Subunit 6 of the COP9 signalosome promotes tumorigenesis in mice through stabilization of MDM2 and is upregulated in human cancers. *J Clin Invest* 121: 851-865.
46. Lee MH, Zhao R, Phan L, Yeung SC (2011) Roles of COP9 signalosome in cancer. *Cell Cycle* 10: 3057-3066.
47. Tomoda K, Kubota Y, Kato J (1999) Degradation of the cyclin-dependent-kinase inhibitor p27Kip1 is instigated by Jab1. *Nature* 398: 160-165.
48. Naumann M, Bech-Otschir D, Huang X, Ferrell K, Dubiel W (1999) COP9 signalosome-directed c-Jun activation/stabilization is independent of JNK. *J Biol Chem* 274: 35297-35300.
49. Bech-Otschir D, Kraft R, Huang X, Henklein P, Kapelari B, et al. (2001) COP9 signalosome-specific phosphorylation targets p53 to degradation by the ubiquitin system. *EMBO J* 20: 1630-1639.
50. Oh W, Lee EW, Sung YH, Yang MR, Ghim J, et al. (2006) Jab1 induces the cytoplasmic localization and degradation of p53 in coordination with Hdm2. *J Biol Chem* 281: 17457-17465.
51. Brady CA, Attardi LD (2010) p53 at a glance. *J Cell Sci* 123: 2527-2532.
52. Smith J, Tho LM, Xu N, Gillespie DA (2010) The ATM-Chk2 and ATR-Chk1 pathways in DNA damage signaling and cancer. *Adv Cancer Res* 108: 73-112.
53. Huang J, Yuan H, Lu C, Liu X, Cao X, et al. (2007) Jab1 mediates protein degradation of the Rad9-Rad1-Hus1 checkpoint complex. *J Mol Biol* 371: 514-527.
54. Yang X, Menon S, Lykke-Andersen K, Tsuge T, Di Xiao, et al. (2002) The COP9 signalosome inhibits p27(kip1) degradation and impedes G1-S phase progression via deneddylation of SCF Cul1. *Curr Biol* 12: 667-672.
55. Denti S, Fernandez-Sanchez ME, Rogge L, Bianchi E (2006) The COP9 signalosome regulates Skp2 levels and proliferation of human cells. *J Biol Chem* 281: 32188-32196.
56. Bondar T, Kalinina A, Khair L, Kopanja D, Nag A, et al. (2006) Cul4A and DDB1 associate with Skp2 to target p27Kip1 for proteolysis involving the COP9 signalosome. *Mol Cell Biol* 26: 2531-2539.
57. Panattoni M, Sanvito F, Basso V, Doglioni C, Casorati G, et al. (2008) Targeted inactivation of the COP9 signalosome impairs multiple stages of T cell development. *J Exp Med* 205: 465-477.
58. Menon S, Chi H, Zhang H, Deng XW, Flavell RA, et al. (2007) COP9 signalosome subunit 8 is essential for peripheral T cell homeostasis and antigen receptor-induced entry into the cell cycle from quiescence. *Nat Immunol* 8: 1236-1245.
59. Kob R, Kelm J, Posorski N, Baniahmad A, von Eggeling F, et al. (2009) Regulation of the anaphase-promoting complex by the COP9 signalosome. *Cell Cycle* 8: 2041-2049.
60. Fingerle-Rowson G, Petrenko O (2007) MIF coordinates the cell cycle with DNA damage checkpoints. Lessons from knockout mouse models. *Cell Div* 2: 22.
61. Lykke-Andersen K, Schaefer L, Menon S, Deng XW, Miller JB, et al. (2003) Disruption of the COP9 signalosome Csn2 subunit in mice causes deficient cell proliferation, accumulation of p53 and cyclin E, and early embryonic death. *Mol Cell Biol* 23: 6790-6797.
62. Yan J, Walz K, Nakamura H, Carattini-Rivera S, Zhao Q, et al. (2003) COP9 signalosome subunit 3 is essential for maintenance of cell proliferation in the mouse embryonic epiblast. *Mol Cell Biol* 23: 6798-6808.
63. Tomoda K, Yoneda-Kato N, Fukumoto A, Yamanaka S, Kato JY (2004) Multiple functions of Jab1 are required for early embryonic development and growth potential in mice. *J Biol Chem* 279: 43013-43018.
64. Orel L, Neumeier H, Hochrainer K, Binder BR, Schmid JA (2010) Crosstalk between the NF-kappaB activating IKK-complex and the CSN signalosome. *J Cell Mol Med* 14: 1555-1568.
65. Karin M (2009) NF-kappaB as a critical link between inflammation and cancer. *Cold Spring Harb Perspect Biol* 1: a000141.
66. Uhle S, Medalia O, Waldron R, Dumdey R, Henklein P, et al. (2003) Protein kinase CK2 and protein kinase D are associated with the COP9 signalosome. *EMBO J* 22: 1302-1312.
67. Henke W, Ferrell K, Bech-Otschir D, Seeger M, Schade R, et al. (1999) Comparison of human COP9 signalosome and 26S proteasome lid'. *Mol Biol Rep* 26: 29-34.
68. Huang MT, Ma W, Lu YP, Chang RL, Fisher C, et al. (1995) Effects of curcumin, demethoxycurcumin, bisdemethoxycurcumin and tetrahydrocurcumin on 12-O-tetradecanoylphorbol-13-acetate-induced tumor promotion. *Carcinogenesis* 16: 2493-2497.
69. Arbiser JL, Klauber N, Rohan R, van Leeuwen R, Huang MT, et al. (1998) Curcumin is an in vivo inhibitor of angiogenesis. *Mol Med* 4: 376-383.
70. Crawford LJ, Walker B, Irvine AE (2011) Proteasome inhibitors in cancer therapy. *J Cell Commun Signal* 5: 101-110.

71. Moreau P, Richardson PG, Cavo M, Orłowski RZ, San Miguel JF, et al. (2012) Proteasome inhibitors in multiple myeloma: 10 years later. *Blood* 120: 947-959.
72. Ferlay J, Shin HR, Bray F, Forman D, Mathers C, et al. (2010) Estimates of worldwide burden of cancer in 2008: GLOBOCAN 2008. *Int J Cancer* 127: 2893-2917.
73. Kirstein MM, Vogel A (2014) The pathogenesis of hepatocellular carcinoma. *Dig Dis* 32: 545-553.
74. Kaposi-Novak P, Libbrecht L, Woo HG, Lee YH, Sears NC, et al. (2009) Central role of c-Myc during malignant conversion in human hepatocarcinogenesis. *Cancer Res* 69: 2775-2782.
75. Li B, Simon MC (2013) Molecular Pathways: Targeting MYC-induced metabolic reprogramming and oncogenic stress in cancer. *Clin Cancer Res* 19: 5835-5841.
76. Panattoni M, Maiorino L, Lukacs A, Zentilin L, Mazza D, et al. (2014) The COP9 signalosome is a repressor of replicative stress responses and polyploidization in the regenerating liver. *Hepatology* 59: 2331-2343.
77. Lee YH, Judge AD, Seo D, Kitade M, Gómez-Quiroz LE, et al. (2011) Molecular targeting of CSN5 in human hepatocellular carcinoma: a mechanism of therapeutic response. *Oncogene* 30: 4175-4184.
78. Koga H, Harada M, Ohtsubo M, Shishido S, Kumemura H, et al. (2003) Troglitazone induces p27Kip1-associated cell-cycle arrest through down-regulating Skp2 in human hepatoma cells. *Hepatology* 37: 1086-1096.
79. Koga H, Sakisaka S, Harada M, Takagi T, Hanada S, et al. (2001) Involvement of p21(WAF1/Cip1), p27(Kip1), and p18(INK4c) in troglitazone-induced cell-cycle arrest in human hepatoma cell lines. *Hepatology* 33: 1087-1097.
80. Wong N, Yeo W, Wong WL, Wong NL, Chan KY, et al. (2009) TOP2A overexpression in hepatocellular carcinoma correlates with early age onset, shorter patients survival and chemoresistance. *Int J Cancer* 124: 644-652.
81. Zhang A, Lyu YL, Lin CP, Zhou N, Azarova AM, et al. (2006) A protease pathway for the repair of topoisomerase II-DNA covalent complexes. *J Biol Chem* 281: 35997-36003.
82. Chen MC, Chen CH, Chuang HC, Kulp SK, Teng CM, et al. (2011) Novel mechanism by which histone deacetylase inhibitors facilitate topoisomerase II alpha degradation in hepatocellular carcinoma cells. *Hepatology* 53: 148-159.
83. Lu YS, Kashida Y, Kulp SK, Wang YC, Wang D, et al. (2007) Efficacy of a novel histone deacetylase inhibitor in murine models of hepatocellular carcinoma. *Hepatology* 46: 1119-1130.
84. Yu YS, Tang ZH, Pan QC, Chen XH, Liu XN, et al. (2012) Inhibition of Csn3 expression induces growth arrest and apoptosis of hepatocellular carcinoma cells. *Cancer Chemother Pharmacol* 69: 1173-1180.
85. Lei D, Li F, Su H, Tian Z, Ye B, et al. (2011) COP9 signalosome subunit 8 is required for postnatal hepatocyte survival and effective proliferation. *Cell Death Differ* 18: 259-270.
86. Tanaka Y, Kanai F, Ichimura T, Tateishi K, Asaoka Y, et al. (2006) The hepatitis B virus X protein enhances AP-1 activation through interaction with Jab1. *Oncogene* 25: 633-642.
87. Parsonnet J, Friedman GD, Vandersteen DP, Chang Y, Vogelman JH, et al. (1991) *Helicobacter pylori* infection and the risk of gastric carcinoma. *N Engl J Med* 325: 1127-1131.
88. Liu N, Liu X, Zhou N, Wu Q, Zhou L, et al. (2014) Gene expression profiling and bioinformatics analysis of gastric carcinoma. *Exp Mol Pathol* 96: 361-366.
89. Oshimo Y, Oue N, Mitani Y, Nakayama H, Kitadai Y, et al. (2004) Frequent loss of RUNX3 expression by promoter hypermethylation in gastric carcinoma. *Pathobiology* 71: 137-143.
90. Ito K, Liu Q, Salto-Tellez M, Yano T, Tada K, et al. (2005) RUNX3, a novel tumor suppressor, is frequently inactivated in gastric cancer by protein mislocalization. *Cancer Res* 65: 7743-7750.
91. Kim JH, Choi JK, Cinghu S, Jang JW, Lee YS, et al. (2009) Jab1/CSN5 induces the cytoplasmic localization and degradation of RUNX3. *J Cell Biochem* 107: 557-565.
92. Crone SG, Jacobsen A, Federspiel B, Bardram L, Krogh A, et al. (2012) microRNA-146a inhibits G protein-coupled receptor-mediated activation of NF- $\kappa$ B by targeting CARD10 and COP8 in gastric cancer. *Mol Cancer* 11: 71.
93. Wu WK, Cho CH, Lee CW, Fan D, Wu K, et al. (2010) Dysregulation of cellular signaling in gastric cancer. *Cancer Lett* 295: 144-153.
94. Murray D, Horgan G, Macmathuna P, Doran P (2008) NET1-mediated RhoA activation facilitates lysophosphatidic acid-induced cell migration and invasion in gastric cancer. *Br J Cancer* 99: 1322-1329.
95. Winston JT, Strack P, Beer-Romero P, Chu CY, Elledge SJ, et al. (1999) The SCFbeta-TRCP-ubiquitin ligase complex associates specifically with phosphorylated destruction motifs in IkappaBalpha and beta-catenin and stimulates IkappaBalpha ubiquitination in vitro. *Genes Dev* 13: 270-283.
96. Shin JY, Kim HS, Lee KS, Kim J, Park JB, et al. (2000) Mutation and expression of the p27KIP1 and p57KIP2 genes in human gastric cancer. *Exp Mol Med* 32: 79-83.
97. Lee MH, Yang HY (2001) Negative regulators of cyclin-dependent kinases and their roles in cancers. *Cell Mol Life Sci* 58: 1907-1922.
98. Lee MH, Reynisdóttir I, Massagué J (1995) Cloning of p57KIP2, a cyclin-dependent kinase inhibitor with unique domain structure and tissue distribution. *Genes Dev* 9: 639-649.
99. Kamura T, Hara T, Kotshiba S, Yada M, Ishida N, et al. (2003) Degradation of p57Kip2 mediated by SCFSkp2-dependent ubiquitylation. *Proc Natl Acad Sci U S A* 100: 10231-10236.
100. Wei Z, Jiang X, Liu F, Qiao H, Zhou B, et al. (2013) Downregulation of Skp2 inhibits the growth and metastasis of gastric cancer cells in vitro and in vivo. *Tumour Biol* 34: 181-192.
101. Chen B, Zhao R, Su CH, Linan M, Tseng C, et al. (2012) CDK inhibitor p57 (Kip2) is negatively regulated by COP9 signalosome subunit 6. *Cell Cycle* 11: 4633-4641.
102. World-Health-Organization (2014) World Cancer Report 2014.
103. Iodice S, Gandini S, Maisonneuve P, Lowenfels AB (2008) Tobacco and the risk of pancreatic cancer: a review and meta-analysis. *Langenbecks Arch Surg* 393: 535-545.
104. Bond-Smith G, Banga N, Hammond TM, Imber CJ (2012) Pancreatic adenocarcinoma. *BMJ* 344: e2476.
105. Kouvaraki MA, Korapati AL, Rassidakis GZ, Tian L, Zhang Q, et al. (2006) Potential role of Jun activation domain-binding protein 1 as a negative regulator of p27kip1 in pancreatic adenocarcinoma. *Cancer Res* 66: 8581-8589.
106. Solinas-Toldo S, Wallrapp C, Müller-Pillasch F, Bentz M, Gress T, et al. (1996) Mapping of chromosomal imbalances in pancreatic carcinoma by comparative genomic hybridization. *Cancer Res* 56: 3803-3807.
107. Tomoda K, Kubota Y, Arata Y, Mori S, Maeda M, et al. (2002) The cytoplasmic shuttling and subsequent degradation of p27Kip1 mediated by Jab1/CSN5 and the COP9 signalosome complex. *J Biol Chem* 277: 2302-2310.
108. Carrano AC, Eytan E, Hershko A, Pagano M (1999) SKP2 is required for ubiquitin-mediated degradation of the CDK inhibitor p27. *Nat Cell Biol* 1: 193-199.
109. Tanaka T, Iino M (2014) Sec6 regulated cytoplasmic translocation and degradation of p27 via interactions with Jab1 and Siah1. *Cell Signal* 26: 2071-2085.
110. Jumpertz S, Hennes T, Asare Y, Vervoorts J, Bernhagen J, et al. (2014) The  $\beta$ -catenin E3 ubiquitin ligase SIAH-1 is regulated by CSN5/JAB1 in CRC cells. *Cell Signal* 26: 2051-2059.
111. Nagano Y, Fukushima T, Okemoto K, Tanaka K, Bowtell DD, et al. (2011) Siah1/SIP regulates p27(kip1) stability and cell migration under metabolic stress. *Cell Cycle* 10: 2592-2602.
112. Malicet C, Hoffmeister A, Moreno S, Closa D, Dagorn JC, et al. (2006) Interaction of the stress protein p8 with Jab1 is required for Jab1-dependent p27 nuclear-to-cytoplasm translocation. *Biochem Biophys Res Commun* 339: 284-289.
113. Su SB, Motoo Y, Iovanna JL, Xie MJ, Mouri H, et al. (2001) Expression of p8 in human pancreatic cancer. *Clin Cancer Res* 7: 309-313.

114. Savkovic V, Gaiser S, Iovanna JL, Bödeker H (2004) The stress response of the exocrine pancreas. *Dig Dis* 22: 239-246.
115. Singh P, Wig JD, Srinivasan R (2011) The Smad family and its role in pancreatic cancer. *Indian J Cancer* 48: 351-360.
116. Wan M, Huang J, Jhala NC, Tytler EM, Yang L, et al. (2005) SCF(beta-TrCP1) controls Smad4 protein stability in pancreatic cancer cells. *Am J Pathol* 166: 1379-1392.
117. Cowan RW, Maitra A (2014) Genetic progression of pancreatic cancer. *Cancer J* 20: 80-84.
118. Wan M, Cao X, Wu Y, Bai S, Wu L, et al. (2002) Jab1 antagonizes TGF-beta signaling by inducing Smad4 degradation. *EMBO Rep* 3: 171-176.
119. Fukumoto A, Tomoda K, Yoneda-Kato N, Nakajima Y, Kato JY (2006) Depletion of Jab1 inhibits proliferation of pancreatic cancer cell lines. *FEBS Lett* 580: 5836-5844.
120. Li J, Wang Y, Yang C, Wang P, Oelschläger DK, et al. (2009) Polyethylene glycosylated curcumin conjugate inhibits pancreatic cancer cell growth through inactivation of Jab1. *Mol Pharmacol* 76: 81-90.
121. Schütz AK, Hennes T, Jumpertz S, Fuchs S, Bernhagen J (2012) Role of CSN5/JAB1 in Wnt/ $\beta$ -catenin activation in colorectal cancer cells. *FEBS Lett* 586: 1645-1651.
122. Phelps RA, Chidester S, Dehghanizadeh S, Phelps J, Sandoval IT, et al. (2009) A two-step model for colon adenoma initiation and progression caused by APC loss. *Cell* 137: 623-634.
123. Kinzler KW, Vogelstein B (1996) Lessons from hereditary colorectal cancer. *Cell* 87: 159-170.
124. Ilyas M, Tomlinson IP, Rowan A, Pignatelli M, Bodmer WF (1997) Beta-catenin mutations in cell lines established from human colorectal cancers. *Proc Natl Acad Sci U S A* 94: 10330-10334.
125. Clevers H, Nusse R (2012) Wnt/ $\beta$ -catenin signaling and disease. *Cell* 149: 1192-1205.
126. Huang X, Langelotz C, Hetfeld-Pechoc BK, Schwenk W, Dubiel W (2009) The COP9 signalosome mediates beta-catenin degradation by deneddylation and blocks adenomatous polyposis coli destruction via USP15. *J Mol Biol* 391: 691-702.
127. Xin D, Rendon BE, Zhao M, Winner M, McGhee Coleman A, et al. (2010) The MIF homologue D-dopachrome tautomerase promotes COX-2 expression through  $\beta$ -catenin-dependent and -independent mechanisms. *Mol Cancer Res* 8: 1601-1609.
128. Steinbach G, Lynch PM, Phillips RK, Wallace MH, Hawk E, et al. (2000) The effect of celecoxib, a cyclooxygenase-2 inhibitor, in familial adenomatous polyposis. *N Engl J Med* 342: 1946-1952.
129. Brown JR, DuBois RN (2005) COX-2: a molecular target for colorectal cancer prevention. *J Clin Oncol* 23: 2840-2855.
130. Matsuzawa SI, Reed JC (2001) Siah-1, SIP, and Ebi collaborate in a novel pathway for beta-catenin degradation linked to p53 responses. *Mol Cell* 7: 915-926.
131. Liu J, Stevens J, Rote CA, Yost HJ, Hu Y, et al. (2001) Siah-1 mediates a novel beta-catenin degradation pathway linking p53 to the adenomatous polyposis coli protein. *Mol Cell* 7: 927-936.
132. Gonzalez-Moreno S, Gonzalez-Bayon LA, Ortega-Perez G (2010) Hyperthermic intraperitoneal chemotherapy: Rationale and technique. *World J Gastrointest Oncol* 2: 68-75.
133. Feist M, Huang X, Müller JM, Rau B, Dubiel W (2014) Can hyperthermic intraperitoneal chemotherapy efficiency be improved by blocking the DNA repair factor COP9 signalosome? *Int J Colorectal Dis* 29: 673-680.
134. Leppert U, Henke W, Huang X, Müller JM, Dubiel W (2011) Post-transcriptional fine-tuning of COP9 signalosome subunit biosynthesis is regulated by the c-Myc/Lin28B/let-7 pathway. *J Mol Biol* 409: 710-721.
135. Loda M, Cukor B, Tam SW, Lavin P, Fiorentino M, et al. (1997) Increased proteasome-dependent degradation of the cyclin-dependent kinase inhibitor p27 in aggressive colorectal carcinomas. *Nat Med* 3: 231-234.