



Supplementary Figure 1: Plot of the mean Aptensio adult (blue line), children 6 to 12 (black line), and children 4 to 5 (red line) years old plasma levels vs. time. Note: (•••••) ADULT; (— — —) CHILD_4TO5; (— — —) CHILD_6TO12.

Correspondence to: Andre J Jackson, Department of Pharmacy, Howard University, 2300 4th Street Washington DC, USA, Tel: +13016468602; E-mail: Jacksonan1945@gmail.com

Received: 04-Jun-2022, Manuscript No. JBB-22-16895; **Editor assigned:** 08-Jun-2022, PreQC No. JBB-22-16895 (PQ); **Reviewed:** 24-Jun-2022, QC No. JBB-22-16895; **Revised:** 01-Jul-2022, Manuscript No. JBB-22-16895 (R); **Published:** 08-Jul-2022, DOI:10.35248/0975-0851.22.14.474.

Citation: Jackson AJ, Brundage R, Ette E, Chaudhary I (2022) Semi-Physiological based Population Pharmacokinetic Model for Methylphenidate Hydrochloride Multi-Layer Extended Release (Aptensio XR[®]) Capsules in Children ages 4-12 and Adults. J Bioequiv Availab. 14:474.

Copyright: © 2022 Jackson AJ, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.